

**INFORMED CONSENT FOR MEDICATION**

**Dosage and / or Side Effect information last revised on 12/17/2010**

Completion of this form is voluntary. If not completed, the medication cannot be administered without a court order unless in an emergency.  
 This consent is maintained in the client's record and is accessible to authorized users.

Name – Patient / Client (Last, First, MI)		ID Number	Living Unit	Birthdate
Name – Individual Preparing This Form		Name – Staff Contact		Name / Telephone Number – Institution

MEDICATION CATEGORY	MEDICATION	RECOMMENDED DAILY TOTAL DOSAGE RANGE	ANTICIPATED DOSAGE RANGE
Antipsychotic Agent	Seroquel; Seroquel XR (quetiapine)	Quetiapine 50mg - 800mg Quetiapine XR 400mg - 800mg	

The anticipated dosage range is to be individualized, may be above or below the recommended range but no medication will be administered without your informed and written consent.

Recommended daily total dosage range of manufacturer, as stated in Physician's Desk Reference (PDR) or another standard reference.

This medication will be administered  Orally  Injection  Other – Specify:

**1. Reason for Use of Psychotropic Medication and Benefits Expected (note if this is 'Off Label' Use)**

Include DSM IV diagnosis or the diagnostic "working hypothesis."

**2. Alternative mode(s) of treatment other than or in addition to medications include**

Note: Some of these would be applicable only in an inpatient environment.

- |  |  |
|--|--|
| <input type="checkbox"/> -Environment and / or staff changes         | <input type="checkbox"/> -Rehabilitation treatments / therapy (OT, PT, AT) |
| <input type="checkbox"/> -Positive redirection and staff interaction | <input type="checkbox"/> -Treatment programs and approaches (habilitation) |
| <input type="checkbox"/> -Individual and / or group therapy          | <input type="checkbox"/> -Use of behavior intervention techniques          |

**Other Alternatives:**

**3. Probable consequences of NOT receiving the proposed medication are**

**Impairment of**  -Work Activities  -Family Relationships  -Social Functioning

**Possible increase in symptoms leading to potential**

- |   |   |
|---|---|
| <input type="checkbox"/> -Use of seclusion or restraints                  | <input type="checkbox"/> -Limits on recreation and leisure activities |
| <input type="checkbox"/> -Limits on access to possessions                 | <input type="checkbox"/> -Intervention of law enforcement authorities |
| <input type="checkbox"/> -Limits on personal freedoms                     | <input type="checkbox"/> -Risk of harm to self or others              |
| <input type="checkbox"/> -Limit participation in treatment and activities |   |

**Other consequences**

**Note:** These consequences may vary, depending upon whether or not the individual is in an inpatient setting. It is also possible that in unusual situations, little or no adverse consequences may occur if the medications are not administered.

Medication : Seroquel;  
Seroquel XR - (quetiapine)

4. Possible side effects, warnings and cautions associated with this medication are listed below. This is not an all inclusive list but is representative of items of potential clinical significance to you. For more information on this medication, you may consult further with your physician or refer to a standard text such as the PDR or the United States Pharmacopoeia Dispensing Information (USPDI). As part of monitoring some of these potential side effects, your physician may order laboratory or other tests. The treatment team will closely monitor individuals who are unable to readily communicate side effects, in order to enhance care and treatment.

Continued – Possible side effects, warnings and cautions associated with this medication.

The most common side effects include: constipation; drowsiness; dry mouth; increased weight; indigestion.

Less common side effects include: abdominal pain; abnormal vision; decrease in appetite; decreased strength and energy; feeling of fast or irregular heartbeat; headache; increased muscle tone; increased sweating; stuffy or runny nose.

Check with your doctor as soon as possible if any of the following side effects occur: dizziness, lightheadedness, or fainting, especially when getting up from a lying or sitting position; fever, chills, muscle aches, or sore throat; loss of balance control; mask-like face; shuffling walk; skin rash; slowed movements; stiffness of arms or legs; swelling of feet or lower legs; trembling and shaking of hands and fingers; trouble in breathing, speaking, or swallowing.

Rare side effects include: fainting; fast, pounding, or irregular heartbeat; menstrual changes; unusual secretion of milk (in females). Symptoms of underactive thyroid--usually two or more occur together--these effects do not require medical attention if they occur alone unless they continue or are bothersome: Dry, puffy skin; loss of appetite; tiredness; weight gain.

**CAUTION** – This medicine may add to the effects of alcohol and other CNS depressants (medicines that make you drowsy or less alert). Check with your doctor before taking any other medications.

Quetiapine may cause drowsiness, especially during the first week of use. Make sure you know how you react to this medicine before you drive, use machines, or do anything else that could be dangerous if you are not alert.

Dizziness, lightheadedness, or fainting may occur, especially when you get up from a lying or sitting position. Getting up slowly may help. If the problem continues or gets worse, check with your doctor.

### **BLACK BOX WARNING**

**Increased Mortality in Elderly Patients with Dementia Related Psychosis:** Elderly patients with dementia related psychosis treated with atypical antipsychotic drugs are at an increased risk of death compared to placebo. Analyses of 17 placebo controlled trials (modal duration of 10 weeks, largely in patients taking atypical antipsychotic drugs, revealed a risk of death in the drug treated patients of between 1.6 to 1.7 times that seen in placebo treated patients. Over the course of a typical 10-week controlled trial, the rate of death in drug treated patients was about 4.5% compared to a rate of about 2.6% in the placebo group. Although the causes of death were varied, most of the deaths appeared to be either cardiovascular (e.g., heart failure, sudden death) or infectious (e.g., pneumonia) in nature. Observational studies suggest that, similar to atypical antipsychotic drugs, treatment with conventional antipsychotic drugs may increase mortality. The extent to which the findings of increased mortality in observational studies may be attributed to the antipsychotic drug as opposed to some characteristic(s) of the patients is not clear.

This drug is not approved for the treatment of patients with dementia-related psychosis.

**Suicidal Ideation in Children, Adolescents, and Young Adults:** Antidepressants increased the risk compared to placebo of suicidal thinking and behavior (suicidality) in short term studies in children, adolescents, and young adults with major depressive disorder (MDD) and other psychiatric disorders. Anyone considering the use of this drug or any other antidepressant in a child, adolescent, or young adult must balance this risk with the clinical need. Short term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction in risk with antidepressants compared to placebo in adults aged 65 and older. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on therapy should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior. Families and care givers should be advised of the need for close observation and communication with the prescriber. This drug is not approved for use in pediatric patients.

See PDR, USPDI or US Hospital Formulary Service for all-inclusive list of side effects.

Client Initial \_\_\_\_\_ Date \_\_\_\_\_

Medication : Seroquel;  
Seroquel XR - (quetiapine)

**By my signature below, I GIVE consent for the named medication on Page 1 and anticipated dosage range. My signature also indicates that I understand the following:**

1. I can refuse to give consent or can withdraw my consent at any time with written notification to the institution director or designee. This will not affect my right to change my decision at a later date. If I withdraw consent after a medication is started, I realize that the medication may not be discontinued immediately. Rather it will be tapered as rapidly as medically safe and then discontinued so as to prevent an adverse medical consequence, such as seizures, due to rapid medication withdrawal.
2. Questions regarding this medication can be discussed with the Interdisciplinary Team, including the physician. The staff contact person can assist in making any necessary arrangements.
3. Questions regarding any behavior support plan or behavior intervention plan, which correspond with the use of the medication, can be directed to the client's social worker, case manager or psychologist.
4. I have the right to request a review at any time of my record, pursuant to ss. 51.30(4)(d) or 51.30(5)(b).
5. I have a legal right to file a complaint if I feel that client rights have been inappropriately restricted. The client's social worker, case manager or agency / facility client rights specialist may be contacted for assistance.
6. My consent permits the dose to be changed within the **anticipated dosage range** without signing another consent.
7. I understand the reasons for the use of the medication, its potential risks and benefits, other alternative treatment(s) and the probable consequences, which may occur if the proposed medication is not given. I have been given adequate time to study the information and find the information to be specific, accurate and complete.
8. This medication consent is for a period effective immediately and not to exceed fifteen (15) months from the date of my signature. The need for and continued use of this medication will be reviewed at least quarterly by the Interdisciplinary Team. The goal, on behalf of the client, will be to arrive at and maintain the client at the minimum effective dose.

**SIGNATURES**

**DATE SIGNED**

Client – If Presumed Competent to Consent/Parent of Minor/Guardian (POA-HC)	Relationship to Client <input type="checkbox"/> Self <input type="checkbox"/> Parent <input type="checkbox"/> Guardian (POA-HC)	
Staff Present at Oral Discussion	Title	

Client / Parent of Minor / Guardian (POA-HC) Comments

**As parent/guardian (POA-HC) was not available for signature, he/she was verbally informed of the information in this consent.**

**Verbal Consent**

Obtained by – PRINT – Staff Name	Date Obtained	Written Consent Received
Obtained from – PRINT – Parent / Guardian (POA-HC) Name	Date Expires	Date Received

Client Initial \_\_\_\_\_ Date \_\_\_\_\_