

COLON PREPS IN CERNER

You need 3 pre-made and saved as favorites “colon preps”. Once you “save” them using the “click button” below the “Power Order Set” you can use them with a single touch. **Please, save instructions to give to all Residents rotating with us.**

1. Activate COLON PREP AM Power Order Set: is meant as a “split prep” starting at 6 PM the day before colonoscopy. Do the following:
 - a) Start PEG (Colyte) at 6 PM,
 - b) Start Dulcolax (Bisacodyl) at 6 PM,
 - c) Start Reglan at 6 PM and repeat in 2 hours if needed,
 - d) Add order “offer Halls mint drops to dissolve in the mouth, to mask colyte flavor while taking the prep”,
 - e) Start Mg Citrate at 4:30 am or 2 1/2 hours before colonoscopy time, whichever is later,
 - f) Be sure there is an order “NPO after midnight, except for medicines and colon prep”,
 - g) Save as **Favorite** under title **“Standard Colon Prep for Next day colonoscopy”**.
2. Activate COLON PREP PM Power Order Set: is meant as a usually “non-split” quick prep when planning to do colonoscopy on the day of the order. Do the following if procedure has not been scheduled and it is earlier than 3 AM; this will allow to do procedure by 7 or 7:30 AM if slot is available; otherwise you can add Mg Citrate 300 mL 2 ½ before scheduled time.
 - a) Start PEG (Colyte) at 3 AM,
 - b) Start Dulcolax (Bisacodyl) at 3 AM,
 - c) Start Reglan at 3 AM and repeat if needed in 2 hours,
 - d)) Add order “offer Halls mint drops to dissolve in the mouth, to mask colyte flavor while taking the prep”,
 - e) Order: “NPO after colon prep except for medicines”,
 - f) Order: Call Endoscopy at 7 AM to find slot time of colonoscopy;
 - g) Order: If colonoscopy will be after 10:30 AM, give 1 bottle of 300 mL of Mg Citrate PO, 2 ½ hours before colonoscopy time,
 - g) Save as **favorite** with name **“Colonoscopy Prep for same day exam; 3 AM start”**
3. Activate COLON PREP PM Power Order Set: is meant as a usually “non-split” quick prep when planning to do colonoscopy on the day of the order. Do the following if procedure has not been scheduled yet and it is later than 3 AM:
 - a) Start PEG Stat,
 - b) Start Dulcolax (Bisacodyl) Stat,
 - c) Start Reglan Stat and repeat if needed in 2 hours,
 - d) Add order “offer Halls mint drops to dissolve in the mouth, to mask colyte flavor while taking the prep”,
 - e) Order: “NPO after colon prep except for medicines”,
 - f) Order: Call Endoscopy at 7 AM or now (whichever is later) to find time of colonoscopy;
 - g) Order: If colonoscopy will be done 5 hours or more after prep is finished, give 1 bottle of 300 mL of Mg Citrate PO, 2 ½ hours before colonoscopy time,
 - h) Save under **Favorites** as **“Colonoscopy Prep for same day exam; STAT start”**