

ORIGINAL ARTICLE

Molecular Architecture of the Goodpasture Autoantigen in Anti-GBM Nephritis

Vadim Pedchenko, Ph.D., Olga Bondar, Ph.D., Agnes B. Fogo, M.D., Roberto Vanacore, Ph.D., Paul Voziyan, Ph.D., A. Richard Kitching, M.D., Ph.D., Jörgen Wieslander, M.D., Ph.D., Clifford Kashtan, M.D., Dorin-Bogdan Borza, Ph.D., Eric G. Neilson, M.D., Curtis B. Wilson, M.D., and Billy G. Hudson, Ph.D.

ABSTRACT

BACKGROUND

In Goodpasture's disease, circulating autoantibodies bind to the noncollagenous-1 (NC1) domain of type IV collagen in the glomerular basement membrane (GBM). The specificity and molecular architecture of epitopes of tissue-bound autoantibodies are unknown. Alport's post-transplantation nephritis, which is mediated by alloantibodies against the GBM, occurs after kidney transplantation in some patients with Alport's syndrome. We compared the conformations of the antibody epitopes in Goodpasture's disease and Alport's post-transplantation nephritis with the intention of finding clues to the pathogenesis of anti-GBM glomerulonephritis.

METHODS

We used an enzyme-linked immunosorbent assay to determine the specificity of circulating autoantibodies and kidney-bound antibodies to NC1 domains. Circulating antibodies were analyzed in 57 patients with Goodpasture's disease, and kidney-bound antibodies were analyzed in 14 patients with Goodpasture's disease and 2 patients with Alport's post-transplantation nephritis. The molecular architecture of key epitope regions was deduced with the use of chimeric molecules and a three-dimensional model of the α 345NC1 hexamer.

RESULTS

In patients with Goodpasture's disease, both autoantibodies to the α 3NC1 monomer and antibodies to the α 5NC1 monomer (and fewer to the α 4NC1 monomer) were bound in the kidneys and lungs, indicating roles for the α 3NC1 and α 5NC1 monomers as autoantigens. High antibody titers at diagnosis of anti-GBM disease were associated with ultimate loss of renal function. The antibodies bound to distinct epitopes encompassing region E_A in the α 5NC1 monomer and regions E_A and E_B in the α 3NC1 monomer, but they did not bind to the native cross-linked α 345NC1 hexamer. In contrast, in patients with Alport's post-transplantation nephritis, alloantibodies bound to the E_A region of the α 5NC1 subunit in the intact hexamer, and binding decreased on dissociation.

CONCLUSIONS

The development of Goodpasture's disease may be considered an autoimmune "conformeropathy" that involves perturbation of the quaternary structure of the α 345NC1 hexamer, inducing a pathogenic conformational change in the α 3NC1 and α 5NC1 subunits, which in turn elicits an autoimmune response. (Funded by the National Institute of Diabetes and Digestive and Kidney Diseases.)

From the Center for Matrix Biology, Division of Nephrology, Department of Medicine (V.P., O.B., A.B.F., R.V., P.V., D.-B.B., E.G.N., B.G.H.), the Departments of Pathology (A.B.F., B.G.H.) and Pediatrics (A.B.F.), and the Department of Biochemistry and the Vanderbilt-Ingram Cancer Center (B.G.H.), Vanderbilt University Medical Center, Nashville; the Center for Inflammatory Diseases, Department of Medicine, Monash University, Clayton, VIC, Australia (A.R.K.); the Department of Nephrology, Lund University Hospital, Lund, Sweden (J.W.); the Department of Pediatrics, University of Minnesota Medical School, Minneapolis (C.K.); and the Department of Immunology and Microbial Science, Scripps Research Institute, La Jolla, CA (C.B.W.). Address reprint requests to Dr. Hudson at the Division of Nephrology, Vanderbilt University Medical Center, S-3223 MCN, 1161 21st Ave. South, Nashville, TN 37232, or at billy.hudson@vanderbilt.edu.

N Engl J Med 2010;363:343-54.

Copyright © 2010 Massachusetts Medical Society.

GOODPASTURE'S DISEASE IS AN ORGAN-specific autoimmune disorder characterized by rapidly progressive glomerulonephritis, pulmonary hemorrhage, and glomerular pathological findings that include linear deposits of antibodies along the glomerular basement membrane (GBM) (Fig. 1A).^{1,2} (For this article we have studied Goodpasture's disease, which describes the specific entity in which the cause of organ dysfunction is proven to be anti-GBM antibodies, in contrast with Goodpasture's syndrome, which is a clinical term used to describe rapidly progressive glomerulonephritis and pulmonary hemorrhage.) Lerner and colleagues³ passively transferred Goodpasture anti-GBM antibodies in a primate model, inducing glomerulonephritis and thereby showing that an autoantibody itself can cause disease. The target GBM antigen for circulating antibodies was subsequently identified as the noncollagenous-1 (NC1) domain of the $\alpha 3$ chain of collagen IV⁴⁻⁶; further studies revealed that collagen IV is a family of six α -chains ($\alpha 1$ through $\alpha 6$).⁷ Immunization of laboratory animals indicated that the $\alpha 3$ NC1 specifically induced severe proteinuria and glomerulonephritis, causally linking the self-antigen and antibody in Goodpasture's disease.⁸⁻¹⁰

The $\alpha 3$ NC1 monomer is assembled into the collagen IV network through the association of the $\alpha 3$, $\alpha 4$, and $\alpha 5$ chains to form a triple helical protomer and through the oligomerization of $\alpha 345$ protomers by means of end-to-end associations and intertwining of triple helices.⁷ Two protomers associate through C-terminal NC1 domains, forming an NC1 hexamer.¹¹ The major cross-linked hexamer is reinforced by novel sulfilimine bonds that fasten two protomers¹² and must be dissociated in order for autoantibody binding to occur.^{11,13} In contrast, the hexamer that is not cross-linked can be dissociated by the antibodies themselves, after which they bind to subunits.¹¹

The $\alpha 345$ network is also a target for anti-GBM alloantibodies in Alport's post-transplantation glomerulonephritis, which occurs in 3 to 5% of patients with Alport's syndrome who receive kidney transplants; in most such patients, the development of Alport's post-transplantation nephritis results in allograft loss.¹⁴ Alport's post-transplantation nephritis is mediated by the deposition of alloantibodies to the $\alpha 3$ NC1 and $\alpha 5$ NC1 domains in response to the "foreign" $\alpha 345$ collagen network that is absent in the kid-

Figure 1 (facing page). Classic Kidney Lesions in Goodpasture's Disease, and the Immunoreactivity of Circulating and Kidney-Bound Goodpasture Autoantibodies to Six Noncollagenous-1 Domain Monomers of Human Collagen IV.

The specimen at left in Panel A (Jones's silver stain) shows cellular crescents (arrows) and necrosis of glomerular tufts (arrowheads), features of glomerulonephritis mediated by anti-glomerular-basement-membrane (GBM) antibodies; the specimen at right shows a glomerulus with crescent and linear staining of the GBM with fluorescein-labeled antihuman IgG antibody. Panels B, C, and D show the reactivity of serum from a total of 57 patients with Goodpasture's disease, grouped according to noncollagenous-1 (NC1) specificity. In Panel B, serum samples from 12 patients react only with $\alpha 3$ NC1. In Panel C, samples from 12 different patients react with $\alpha 3$ NC1 and $\alpha 5$ NC1. In Panel D, samples from 33 different patients react with $\alpha 1$ NC1, $\alpha 3$ NC1, $\alpha 4$ NC1, and $\alpha 5$ NC1. These findings differed significantly from the findings in serum samples from 18 healthy volunteers, which showed nonreactivity ($P < 0.05$). Panel E shows the binding of autoantibodies eluted from the kidneys of 14 patients with Goodpasture's disease. Significant binding was detected only to the $\alpha 3$ NC1, $\alpha 5$ NC1, and $\alpha 4$ NC1 domains, with less binding to the last than to the first two ($P < 0.05$). Normal kidney eluates from 3 patients without Goodpasture's disease were nonreactive with all NC1 domains. In Panels B through E, the circles indicate values in individual patients, the solid horizontal lines indicate medians, and the dotted horizontal lines indicate means plus 3 SD for normal samples.


neys of patients with Alport's syndrome but present in the renal allograft.^{15,16}

Thus, the $\alpha 345$ NC1 hexamer is targeted by antibodies that arise in both Goodpasture's disease and Alport's post-transplantation nephritis, but these antibodies have different binding properties. Alloantibodies bind epitopes exposed on the native hexamer, whereas in Goodpasture's disease the autoantibodies require hexamer dissociation to unmask hidden epitopes.^{7,17} Our retrospective study investigated the molecular basis for these differences in antibody binding to provide insight into the pathogenic mechanisms of autoimmunity in Goodpasture's disease.

METHODS

PROTEINS

We purified recombinant human monomers $\alpha 1$ NC1 through $\alpha 6$ NC1 and chimeras from the culture medium of stably transfected human embryonic kidney (HEK) 293 cells with the use of anti-FLAG agarose.¹⁰ To construct $\alpha 5/\alpha 1$ chime-



ras corresponding to the E_A and E_B regions of the $\alpha 3$ NC1 domain, we used polymerase-chain-reaction mutagenesis (for details see the Supplementary Appendix, available with the full text of this article at NEJM.org). Collagen IV NC1 hexamers were isolated from bovine GBM with the use of collagenase digestion.¹³

SERUM AND TISSUE SAMPLES

Approval from local institutional ethics committees and written informed consent from patients were obtained before the collection of samples. Serum samples from 35 patients with anti-GBM glomerulonephritis were obtained from the serum bank of the Department of Nephrology at

Lund University Hospital as a representative subgroup of samples from a larger cohort that were used in our previous study.¹⁸ An additional 22 serum samples were collected at the Scripps Research Institute, Kansas University, and the Vanderbilt University Medical Center from 1985 through 2008. Samples were collected before plasma exchange or immunosuppressive drug treatment was initiated. Serum samples from 18 healthy adult volunteers were used as normal controls. Tissue eluates were isolated from the kidneys of 13 patients with Goodpasture's disease after they underwent nephrectomy at the Scripps Research Institute, as previously described.^{3,19} Serum and tissue samples obtained at the time of autopsy from one patient with anti-GBM glomerulonephritis who had undergone hemodialysis and immunosuppressive therapy for 3 months²⁰ were snap-frozen, stored at -80°C , and processed later for elution of kidney- and lung-bound antibodies.

Alloantibodies were purified from the rejected kidney allografts of two previously described patients with X-linked Alport's post-transplantation nephritis. Patient 1 was a 23-year-old man with renal insufficiency, proteinuria, and microscopic hematuria.¹⁵ Nephrectomy was performed on a second transplant after linear IgG staining of GBM and crescentic glomerulonephritis were revealed on renal biopsy. In Patient 2, end-stage kidney disease developed at 20 years of age; alloantibodies were eluted from the fourth allograft.¹⁶ Kidneys and lungs from normal donors were obtained from the National Disease Research Interchange in Philadelphia.

Tissue-bound antibodies were eluted with the use of 0.1 M glycine, pH 2.8 and 2.2, after homogenization in TRIS-buffered saline (pH 7.4) with protease inhibitors.²¹

AFFINITY PURIFICATION OF GOODPASTURE AUTOANTIBODIES

The recombinant domain $\alpha 3\text{NC1}$ or $\alpha 5\text{NC1}$ was coupled with Affi-Gel 10 (Bio-Rad Laboratories) at a concentration of 1 mg per milliliter.²² Plasma-pheresis fluid from patients with Goodpasture's disease was fractionated by means of sequential passing through $\alpha 3\text{NC1}$ and $\alpha 5\text{NC1}$ columns. Bound antibodies were eluted with 6 M urea in 50 mM sodium citrate (pH 4.0) diluted with TRIS-buffered saline (pH 7.4) and concentrated with the use of ultrafiltration.

IMMUNOASSAYS

Immunoassays of NC1 domains or chimeras were performed with the use of indirect and inhibition enzyme-linked immunosorbent assays.²³

STATISTICAL ANALYSIS

All data sets were analyzed for normality with the use of the Kolmogorov–Smirnov test. To determine differences between groups, we used the Mann–Whitney U test or the Kruskal–Wallis analysis of variance on ranks for continuous variables and Fisher's exact test for categorical variables. A P value of less than 0.05 was considered to indicate statistical significance.

RESULTS

CLINICAL DATA

For this retrospective study, we included serum samples from 57 patients with Goodpasture's disease. The median age of the patients at the time of diagnosis was 59 years (range, 19 to 87); 44% of all patients were women. There was no significant difference in age distribution between male and female patients. In 3 patients, no further clinical data were available. Among the remaining 54 patients, 22 (41%) had positive results for myeloperoxidase antineutrophil cytoplasmic antibodies (ANCA). Clinical data on lung involvement were available for 46 patients, and 12 of these patients (26%) had overt lung hemorrhage. Follow-up information was available for 50 of the 57 patients at 6 months; 17 patients (34%) remained alive, with stable kidney function; 21 (42%) were being treated with dialysis; and 12 (24%) had died.

SPECIFICITY OF CIRCULATING AND KIDNEY-BOUND ANTIBODIES

Serum samples from all 57 patients with Goodpasture's disease reacted strongly with the $\alpha 3\text{NC1}$ domain. There were three categories of specificity: 12 samples reacted only with the $\alpha 3\text{NC1}$ monomer (Fig. 1B), 12 reacted with both the $\alpha 3\text{NC1}$ and $\alpha 5\text{NC1}$ monomers (Fig. 1C), and the remaining 33 samples were immunoreactive to $\alpha 3\text{NC1}$, $\alpha 5\text{NC1}$, $\alpha 1\text{NC1}$, and $\alpha 4\text{NC1}$, with occasional binding to $\alpha 2\text{NC1}$ and $\alpha 6\text{NC1}$ (Fig. 1D). Overall, 72% of the samples from these patients reacted with the $\alpha 5\text{NC1}$ monomer. The antibodies eluted from the kidneys of all 14 patients with Goodpasture's disease showed binding to $\alpha 3\text{NC1}$ and $\alpha 5\text{NC1}$ monomers in the majority of samples (11 of 14,




Figure 2. Characterization and Epitope Mapping of Circulating Goodpasture Autoantibodies Specific to the $\alpha 3$ and $\alpha 5$ Noncollagenous-1 Domains.

Autoantibodies were preincubated with various concentrations of the monomer $\alpha 3$ or $\alpha 5$ noncollagenous-1 (NC1) domain, and binding to immobilized antigens $\alpha 3$ NC1 and $\alpha 5$ NC1 was measured with the use of an enzyme-linked immunosorbent assay (ELISA). Panels A and B show means (\pm SE) for relative binding, expressed as a percentage of binding in the absence of NC1 monomers in solution, for $\alpha 3$ NC1 and $\alpha 5$ NC1 IgG antibodies, respectively, from seven patients with Goodpasture's disease. Binding of the $\alpha 3$ NC1 IgG antibodies to immobilized $\alpha 3$ NC1 was strongly inhibited in the presence of soluble $\alpha 3$ NC1 (solid circles) but not $\alpha 5$ NC1 (open circles) (Panel A). The $\alpha 5$ NC1 IgG antibodies had a lower affinity for $\alpha 5$ NC1 (Panel B). Panel C shows the extent of binding of $\alpha 3$ NC1 and $\alpha 5$ NC1 IgG antibodies to NC1 hexamers from native glomerular basement membrane (N-GBM) and dissociated GBM (D-GBM). IgG antibodies from individual serum samples from patients with Goodpasture's disease are represented by circles and medians by horizontal lines. Panel D shows the alignment of the $\alpha 1$ NC1 and $\alpha 5$ NC1 amino acid sequences corresponding to the E_A and E_B regions of the $\alpha 3$ NC1 domain. Residues that differ from those in $\alpha 1$ NC1 (bold) and residues that were mutated in $\alpha 5$ chimeras (bold red) are shown. Panel E shows means (\pm SE) for the inhibition of the binding of circulating $\alpha 5$ NC1-IgG antibodies from the seven patients with Goodpasture's disease to the $\alpha 5$ NC1 domain. E_A - $\alpha 5$ chimeras are represented by solid triangles, and E_B - $\alpha 5$ chimeras by open triangles. The monomers $\alpha 5$ NC1 (open circles) and $\alpha 1$ NC1 (solid circles) were included as positive and negative controls, respectively. In Panels A, B, and E, I bars denote standard errors for seven $\alpha 5$ NC1 antibodies.

or 79%) (Fig. 1E), with significantly lower binding to $\alpha 4$ NC1.

CHARACTERIZATION OF CIRCULATING $\alpha 3$ NC1 AND $\alpha 5$ NC1 AUTOANTIBODIES

We purified antibodies from seven patients with Goodpasture's disease, using $\alpha 3$ NC1 and $\alpha 5$ NC1

affinity columns. All purified antibodies belonged to the IgG subclass (data not shown). Binding of the $\alpha 3$ NC1 antibodies to immobilized $\alpha 3$ NC1 was strongly inhibited with soluble $\alpha 3$ NC1 but not with the $\alpha 5$ NC1 monomer (Fig. 2A). Potent $\alpha 3$ NC1 inhibition (half-maximal inhibitory concentration [IC_{50}], 0.05 μ g per milliliter) indi-

cates high affinity of $\alpha 3\text{NC1}$ antibodies (apparent dissociation constant [K_D], 2×10^{-9} M). The $\alpha 5\text{NC1}$ IgG antibodies had lower affinity for the $\alpha 5\text{NC1}$ monomer (IC_{50} , $1.3 \mu\text{g}$ per milliliter; apparent K_D , 5×10^{-8} M) (Fig. 2B). The absence of cross-inhibitory effects of $\alpha 5\text{NC1}$ and $\alpha 3\text{NC1}$ antibodies shows that $\alpha 3\text{NC1}$ antibodies and $\alpha 5\text{NC1}$ antibodies are two distinct populations of circulating autoantibodies in Goodpasture's disease.

Reduction of the $\alpha 5\text{NC1}$ monomer completely inhibited binding of the purified $\alpha 5\text{NC1}$ antibodies (data not shown), indicating that the epitopes are conformational and dependent on a critical disulfide bond, analogous to that of $\alpha 3\text{NC1}$.¹³ Moreover, the $\alpha 3\text{NC1}$ and $\alpha 5\text{NC1}$ antibodies displayed negligible binding to native GBM NC1 hexamers, but the binding was greatly increased on dissociation of the hexamers into constituent subunits (Fig. 2C). We previously described this phenomenon for $\alpha 3\text{NC1}$ antibodies as cryptic (hidden) epitopes.^{13,22}

EPITOPE MAPPING FOR CIRCULATING $\alpha 5\text{NC1}$ GOODPASTURE ANTIBODIES

We hypothesized that regions in the $\alpha 5\text{NC1}$ monomer that were homologous to the E_A and E_B regions of the $\alpha 3\text{NC1}$ monomer^{23,24} would harbor the epitopes for the $\alpha 5\text{NC1}$ antibodies. We created two $\alpha 1/\alpha 5$ chimeras by substituting unique amino acid residues in $\alpha 1\text{NC1}$, as a nonreactive scaffold, for those in $\alpha 5\text{NC1}$ (Fig. 2D). Preincubation with the E_A - $\alpha 5$ chimera, but not with the E_B - $\alpha 5$ chimera or a parental $\alpha 1\text{NC1}$ monomer, significantly inhibited binding of Goodpasture $\alpha 5\text{NC1}$ antibodies to $\alpha 5\text{NC1}$ in a dose-dependent manner (Fig. 2E). These results establish the E_A region as a part of the epitope for circulating $\alpha 5\text{NC1}$ autoantibodies.

EPITOPE MAPPING FOR KIDNEY-BOUND AUTOANTIBODIES AND ALLOANTIBODIES


Both the E_A and E_B regions of the $\alpha 3\text{NC1}$ monomer were targets for kidney-bound antibodies in 11 patients with Goodpasture's disease (Fig. 3A). All kidney eluates also targeted the E_A region of the $\alpha 5\text{NC1}$ monomer, whereas only 1 patient had antibodies that were reactive to the E_B region (Fig. 3B). Moreover, comparison of samples from a single patient with Goodpasture's disease revealed that circulating antibodies and lung-bound and kidney-bound antibodies shared the same specificity, affinity, and epitopes (Fig. 1 in the Supplementary Appendix). In contrast, the allo-

Figure 3 (facing page). Comparison of Kidney- and Lung-Bound Autoantibodies from Patients with Goodpasture's Disease and Alloantibodies from Patients with Alport's Post-Transplantation Nephritis.

Panels A and B show the extent to which kidney-bound Goodpasture autoantibodies bind to the E_A and E_B chimeras of the $\alpha 3$ noncollagenous-1 (NC1) and $\alpha 5\text{NC1}$ domains. Individual patients with Goodpasture's disease are represented by circles, background binding to $\alpha 1\text{NC1}$ by dotted lines, and median values for groups that are different from the background by horizontal lines ($P < 0.05$). Panel C shows the specificity of kidney-bound alloantibodies for the $\alpha 3/\alpha 5\text{NC1}$ monomer and epitope in samples from two patients with Alport's post-transplantation nephritis (APTN). Panel D shows the binding of circulating, kidney-bound, and lung-bound autoantibodies to native glomerular basement membrane (N-GBM) and dissociated GBM (D-GBM) NC1 hexamers in samples from one patient with Goodpasture's disease and two patients with APTN. Normal human IgG (hIgG) does not bind to NC1 hexamers. Panel E shows the binding of circulating antibodies from 27 patients with Goodpasture's disease and kidney-bound autoantibodies from 14 patients with Goodpasture's disease to N-GBM and D-GBM NC1 hexamers. Individual patients are represented by circles, and medians for each group by horizontal lines. Panel F shows the positive correlation between the immunoreactivity of the $\alpha 3\text{NC1}$ and $\alpha 5\text{NC1}$ monomers as revealed by simultaneous enzyme-linked immunosorbent assay (ELISA) for all 57 patients with Goodpasture's disease (Spearman's correlation coefficient, 0.852; $P < 0.001$). Data points representing individual patients are fitted to the exponential curve. Panel G shows the levels of $\alpha 3\text{NC1}$ (triangles) and $\alpha 5\text{NC1}$ (circles) autoantibodies in serum from 17 patients with Goodpasture's disease who had functioning native kidneys and 21 patients who were dependent on dialysis at 6-month follow-up. P values are based on the Mann-Whitney U test.

antibodies in kidney eluates from the two patients with Alport's post-transplantation nephritis (Patients 1 and 2) targeted the $\alpha 5\text{NC1}$ monomer but not the homologous $\alpha 1\text{NC1}$ or $\alpha 3\text{NC1}$ monomer (Fig. 3C), and both strongly bound the E_A - $\alpha 5$ chimera, whereas the E_B - $\alpha 5$ chimera reacted with alloantibodies from Patient 2. These unexpected findings indicate that the E_A region of the $\alpha 5\text{NC1}$ monomer is a critical part of the epitopes in both Goodpasture's disease and Alport's post-transplantation nephritis.

Furthermore, both the $\alpha 3\text{NC1}$ and $\alpha 5\text{NC1}$ autoantibodies were nonreactive to the normal $\alpha 345\text{NC1}$ hexamer until the hexamer was dissociated with protein denaturant. The induction of binding was observed for affinity-purified $\alpha 3\text{NC1}$ and $\alpha 5\text{NC1}$ antibodies from a single patient with Goodpasture's disease (Fig. 3D), circulating antibodies from 27 patients with Good-



pasture's disease, and kidney eluates from 14 other patients with Goodpasture's disease (Fig. 3E). Collectively, these findings indicate that circulating and tissue-bound $\alpha 3NC1$ and $\alpha 5NC1$ antibodies in Goodpasture's disease have identi-

cal properties — that is, their respective epitopes arise only after the dissociation of the NC1 hexamer. In sharp contrast, the alloantibodies associated with Alport's post-transplantation nephritis have a strong reaction to the normal

hexamer, and binding is greatly decreased on dissociation of the hexamer (Fig. 3D).

ASSOCIATION OF α 3NC1 AND α 5NC1 AUTOANTIBODIES WITH DISEASE ACTIVITY

A strong positive correlation was found between titers for α 3NC1 and α 5NC1 antibodies among all serum samples from patients with Goodpasture's disease (Fig. 3F). The presence of the α 3NC1 antibodies in all samples and the gradual increase in α 5NC1 reactivity suggest that α 5NC1 autoantibodies may develop after α 3NC1 autoantibodies.

Further analyses revealed no significant difference in age, sex, ANCA status, renal outcome, or serum reactivity to α 3NC1 or α 5NC1 monomers in patients with and those without lung hemorrhage. ANCA status was not associated with sex, presence or absence of lung involvement, renal outcome, or titers for α 3NC1 and α 5NC1 antibodies; however, patients with positive test results for ANCA were older than patients with negative test results (median age, 70 years vs. 58 years; $P=0.03$). Patients with Goodpasture's disease who were undergoing dialysis and those with preserved renal function at follow-up were of similar age (median, 45 years and 57 years, respectively), but patients who died were significantly older (median, 73 years; $P<0.001$) and were excluded from further analyses. Patients undergoing dialysis had higher titers of α 3NC1 antibodies at presentation than did patients with stable kidney function (Fig. 3G), and had much higher titers for α 5NC1 antibodies (median, 0.922 vs. 0.262). A serum sample from 1 of 21 patients with progressive disease requiring dialysis had reactivity that was restricted to the α 3NC1 monomer; the majority of samples (from 20 of 21 patients) were reactive with α 3NC1 and α 5NC1 monomers. In contrast, samples from 6 of 17 patients with preserved renal function had restricted α 3NC1 reactivity ($P=0.03$ by Fisher's exact test). Thus, our results support the possibility that increased titers of circulating α 3NC1 and α 5NC1 autoantibodies are associated with a poor renal outcome.


THREE-DIMENSIONAL STRUCTURE OF THE α 345NC1 HEXAMER

We analyzed the structure of the immunoreactive E_A and E_B regions in the α 345NC1 hexamer model (see the Supplementary Appendix). The E_A re-

Figure 4 (facing page). Topology of the E_A and E_B Regions in the α 345 Noncollagenous-1 Hexamer, Structural Determinants for the Binding of Alport Alloantibodies and Goodpasture Autoantibodies In Vitro, and Accessible Surface Area of the E_A - α 3 and E_A - α 5 Regions of the Noncollagenous-1 Hexamer.

The α 345 noncollagenous-1 (NC1) hexamer is composed of two trimeric caps, each consisting of α 3NC1 (red), α 4NC1 (blue), and α 5NC1 (green) subunits (Panel A). Two of the six sulfilimine bonds (S=N) that stabilize the trimer-trimer interface are shown (light yellow). The location and structure of the four homologous regions are also shown: E_A (yellow) and E_B (orange) in the α 3NC1 subunit, and E_A (pink) and E_B (purple) in the α 5NC1 subunit. Three regions, E_A and E_B in α 3NC1 and E_A in α 5NC1, become critical parts of the neoepitopes for Goodpasture autoantibodies. The topology of the E_A regions in α 3NC1 and α 5NC1 is similar, as indicated in the ribbon diagrams (Panel A, bottom), with the characteristic folding pattern of a β -sheet stabilized with a disulfide bond. Ala¹⁹, Gln²⁴, and Gln²⁸ (pink) within the E_A region of α 5NC1, exposed in the α 345NC1 hexamer, are candidates for the binding of Alport alloantibodies (Panel B, bottom right). In contrast, Leu²⁷ and Val²⁹ (gray) are sequestered by their lateral interaction with the α 4NC1 domain, and when exposed as a result of hexamer dissociation, they become critical to the binding of Goodpasture autoantibodies. Dissociation of the sulfilimine-cross-linked hexamer into α 35 dimer subunits is concomitant with a conformational change that results in the formation of the neoepitopes encompassing the E_A regions of the α 5NC1 and α 3NC1 monomers and the binding of their respective autoantibodies (Panel B, bottom left). The accessible surface area of the E_A - α 3 region (Panel C, top) and the E_A - α 5 region (Panel C, bottom) was calculated for a probe, which mimics the antibody molecule (radius, 9 Å); the area of individual residues in the α 345NC1 hexamer (black bars) and the α 3NC1/ α 5NC1 model monomers (gray bars) is shown. An increase in the surface area of the monomers indicates that residues are buried in the hexamer (Val²⁷ and Leu²⁹ in E_A - α 3 and Leu²⁷ and Val²⁹ in E_A - α 5). In contrast, residues with similar areas within the hexamer and monomers are exposed in the hexamer (Ala¹⁹, Gln²⁴, and Gln²⁸ in E_A - α 5).

gion of the α 5NC1 subunit was not reactive to the Goodpasture autoantibodies in the α 345NC1 hexamer cross-linked by sulfilimine bonds (Fig. 4A). This lack of reactivity is analogous to that of the E_A and E_B regions of the α 3NC1 subunit.^{11,25} However, disruption of the hexamer quaternary structure after treatment with guanidine or by lowering pH leads to dissociation into α 35 and α 44 dimers and antibody binding (Fig. 4B). The dissociation is concomitant with conformational changes that unlock domain-swapping interactions²⁶



and expose residues sequestered by neighboring subunits. The dissociation and conformational change are reversible, since Goodpasture antibodies do not bind the reassembled hexamer.^{11,25}

Further evidence of conformational transition as a key step in neoepitope formation is provided by the differential effect of dissociating agents on the binding of Goodpasture and Alport post-transplantation nephritis antibodies to the E_A region of the α 5NC1 subunit. Goodpasture autoantibodies react only with the subunits of a dissociated hexamer, whereas Alport post-transplantation nephritis alloantibodies bind to the intact hexamer and lose binding on dissociation. Analysis of the accessible surface area of the E_A- α 5 residues within the α 345 hexamer and in an α 5NC1 monomer reveals that exposure of buried amino acid residues Leu²⁷ and Val²⁹ on hexamer dissociation transforms the E_A- α 5 region into a part of the Goodpasture neoepitope; likewise, homologous residues Val²⁷ and Leu²⁹ become exposed within the E_A- α 3 region (Fig. 4B and 4C). In contrast, Ala¹⁹, Gln²⁴, and Gln²⁸ are located on the hexamer surface and constitute a part of the alloepitope. The diminished binding of the alloantibodies indicates a conformational change in the E_A- α 5 region, which is concomitant with hexamer dissociation.

DISCUSSION

The immunoreactivity of circulating Goodpasture autoantibodies to several NC1 domains of collagen IV was reported previously,²⁷⁻³⁰ but the specificity of tissue-bound autoantibodies is unknown, except in a single patient, in whom the antibodies were reactive to the α 3NC1 domain.³¹ We report here that α 5NC1 autoantibodies, in addition to α 3NC1 autoantibodies, are frequently present in the kidneys and lungs of patients with Goodpasture's disease. The α 5NC1 Goodpasture antibodies bind to a conformation-dependent epitope encompassing the E_A region in the α 5NC1 monomer. This region also encompasses the epitope for alloantibodies in patients with Alport's post-transplantation nephritis.

In the α 345NC1 hexamer, quaternary interactions reinforced by sulfilimine cross-links present key structural constraints against the transition of E_A- α 3 and E_A- α 5 regions to pathogenic conformation in Goodpasture's disease. Disrup-

tion of hexamer structure changes the conformation of the E_A regions of α 3NC1 and α 5NC1 and the E_B region of α 3NC1, transforming them into neoepitopes for autoantibodies. In the GBM, an additional level of constraint is provided by the triple helical domain tethered to the hexamer (conformer 1) (Fig. 5). In the absence of cross-links, quaternary constraints against conformational transition are diminished (conformer 2), shifting the equilibrium toward the trimers (conformer 3). The presence of such trimers in basement membranes has been confirmed on electron microscopy.³² Moreover, Goodpasture antibodies can induce a conformational change, dissociate conformer 3, and form an antigen-antibody complex that is consistent with binding to a non-cross-linked hexamer *in vitro*¹¹ and in passive-transfer experiments.³

We postulate that an early pivotal step of Goodpasture's disease involves conformational transitions in subunits of non-cross-linked hexamers or trimers (conformers 2 and 3), forming pathogenic neoepitopes that elicit both antibody production and binding (conformer 4). The triggering event may be an individual factor or a combination of factors — such as enzymatic or nonenzymatic post-translational modifications (oxidation, nitrosylation, and glycation), a rise in body temperature, or proteolytic cleavage — that perturbs the quaternary structure of the hexamer. Indeed, cleavage of a disulfide bond in α 3NC1 in a non-cross-linked hexamer (conformer 3) has been shown to enhance the binding of Goodpasture antibodies.³³ Furthermore, environmental factors such as cigarette smoking or exposure to organic solvents could inhibit the putative enzyme that catalyzes formation of sulfilimine bonds and thereby increase the proportion of non-cross-linked hexamers (conformer 2).

Goodpasture's disease may be considered an autoimmune "conformeropathy," a designation that reflects the requirement for a conformational transition between two distinct NC1 conformers — a nonpathogenic conformer within the hexamer and a dissociated pathogenic conformer that elicits an autoimmune response. Grave's disease and antiphospholipid autoimmune disease,³⁴⁻³⁸ which involve pathogenic conformational changes, and perhaps idiopathic membranous nephropathy³⁹ may also be included in such a category. This conceptual framework re-




Figure 5. Conformational Diversity and Differential Reactivity of α 345 Noncollagenous-1 Hexamers of the Glomerular Basement Membrane.

The diagram shows a portion of the collagen IV network with the α 345 noncollagenous-1 (NC1) hexamer tethered to the triple-helical domain. The different possible NC1 conformers shown are the cross-linked form stabilized by sulfinimine bonds (conformer 1 [C-1]), the non-cross-linked form (C-2), and the form in which the NC1 hexamers are dissociated into trimers (C-3). In Goodpasture's disease the latter may undergo a conformational change resulting in the formation of neopeptides shown as white squares on the α 3NC1 (red) and α 5NC1 (green) subunits of C-4, eliciting antibody formation and subsequent binding to conformers C-3 and C-4. Conformers C-1 and C-2 have the potential to be transformed into the pathogenic conformer C-4.

flects fundamental issues about the causes of autoimmune disease in molecular terms, answering questions about what triggers the conformational change.

Supported by a grant (DK18381-37) from the National Institute of Diabetes and Digestive and Kidney Diseases (to Dr. Hudson).

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

We thank Parvin Todd and Neonila Danylevych for their technical assistance, Drs. Julia Lewis and Ashton Byington for providing kidney and lung specimens from a patient with Goodpasture's disease, Dr. Julie K. Hudson for critical reading of an earlier version of the manuscript, and Dr. Richard A. Lerner for coining the term "conformeropathy."

REFERENCES

- Wilson CB, Dixon FJ. Anti-glomerular basement membrane antibody-induced glomerulonephritis. *Kidney Int* 1973;3:74-89.
- Salama AD, Levy JB, Lightstone L, Pusey CD. Goodpasture's disease. *Lancet* 2001;358:917-20.
- Lerner RA, Glasscock RJ, Dixon FJ. The role of anti-glomerular basement membrane antibody in the pathogenesis of human glomerulonephritis. *J Exp Med* 1967;126:989-1004.
- Saus J, Wieslander J, Langeveld JP, Quinones S, Hudson BG. Identification of the Goodpasture antigen as the alpha 3(IV) chain of collagen IV. *J Biol Chem* 1988;263:13374-80.
- Butkowski RJ, Langeveld JP, Wieslander J, Hamilton J, Hudson BG. Localization of the Goodpasture epitope to a novel chain of basement membrane collagen. *J Biol Chem* 1987;262:7874-7.
- Turner N, Mason PJ, Brown R, et al. Molecular cloning of the human Goodpasture antigen demonstrates it to be the alpha 3 chain of type IV collagen. *J Clin Invest* 1992;89:592-601.
- Hudson BG, Tryggvason K, Sundaramoorthy M, Neilson EG. Alport's syndrome, Goodpasture's syndrome, and type IV collagen. *N Engl J Med* 2003;348:2543-56.
- Kalluri R, Gattone VH II, Noelken ME, Hudson BG. The alpha 3 chain of type IV collagen induces autoimmune Goodpasture syndrome. *Proc Natl Acad Sci U S A* 1994;91:6201-5.
- Abbate M, Kalluri R, Corna D, et al. Experimental Goodpasture's syndrome in Wistar-Kyoto rats immunized with alpha3 chain of type IV collagen. *Kidney Int* 1998;54:1550-61.
- Sado Y, Boutaud A, Kagawa M, Naito I, Ninomiya Y, Hudson BG. Induction of anti-GBM nephritis in rats by recombinant α 3(IV)NC1 and α 4(IV)NC1 of type IV collagen. *Kidney Int* 1998;53:664-71. [Erratum, *Kidney Int* 1998;54:311.]
- Borza DB, Bondar O, Colon S, et al. Goodpasture autoantibodies unmask cryptic epitopes by selectively dissociating

- autoantigen complexes lacking structural reinforcement: novel mechanisms for immune privilege and autoimmune pathogenesis. *J Biol Chem* 2005;280:27147-54.
12. Vanacore R, Ham AJ, Voehler M, et al. A sulfilimine bond identified in collagen IV. *Science* 2009;325:1230-4.
13. Wieslander J, Langeveld J, Butkowski R, Jodlowski M, Noelken M, Hudson BG. Physical and immunochemical studies of the globular domain of type IV collagen: cryptic properties of the Goodpasture antigen. *J Biol Chem* 1985;260:8564-70.
14. Kashtan CE. Renal transplantation in patients with Alport syndrome. *Pediatr Transplant* 2006;10:651-7.
15. Hudson BG, Kalluri R, Gunwar S, et al. The pathogenesis of Alport syndrome involves type IV collagen molecules containing the alpha 3(IV) chain: evidence from anti-GBM nephritis after renal transplantation. *Kidney Int* 1992;42:179-87.
16. Kang JS, Kashtan CE, Turner AN, Heidet L, Hudson BG, Borza DB. The allotypic sites of alpha3alpha4alpha5(IV) collagen: pathogenic X-linked Alport antibodies target two accessible conformational epitopes in the alpha5NC1 domain. *J Biol Chem* 2007;282:10670-7.
17. Kalluri R, Sun MJ, Hudson BG, Neilson EG. The Goodpasture autoantigen: structural delineation of two immunologically privileged epitopes on alpha3(IV) chain of type IV collagen. *J Biol Chem* 1996;271:9062-8.
18. Segelmark M, Hellmark T, Wieslander J. The prognostic significance in Goodpasture's disease of specificity, titre and affinity of anti-glomerular-basement-membrane antibodies. *Nephron Clin Pract* 2003;94:c59-c68.
19. Marquardt H, Wilson CB, Dixon EJ. Isolation and immunological characterization of human glomerular basement membrane antigens. *Kidney Int* 1973;3:57-65.
20. Clyne S, Frederick C, Arndt F, Lewis J, Fogo AB. Concurrent and discrete clinicopathological presentations of Wegener granulomatosis and anti-glomerular basement membrane disease. *Am J Kidney Dis* 2009;54:1116-20.
21. Saxena R, Bygren P, Butkowski R, Wieslander J. Specificity of kidney-bound antibodies in Goodpasture's syndrome. *Clin Exp Immunol* 1989;78:31-6.
22. Borza DB, Netzer KO, Leinonen A, et al. The Goodpasture autoantigen: identification of multiple cryptic epitopes on the NC1 domain of the alpha3(IV) collagen chain. *J Biol Chem* 2000;275:6030-7.
23. Netzer KO, Leinonen A, Boutaud A, et al. The Goodpasture autoantigen: mapping the major conformational epitope(s) of alpha3(IV) collagen to residues 17-31 and 127-141 of the NC1 domain. *J Biol Chem* 1999;274:11267-74.
24. Hellmark T, Burkhardt H, Wieslander J. Goodpasture disease: characterization of a single conformational epitope as the target of pathogenic autoantibodies. *J Biol Chem* 1999;274:25862-8.
25. Vanacore RM, Ham AJ, Cartailier JP, et al. A role for collagen IV cross-links in conferring immune privilege to the Goodpasture autoantigen: structural basis for the crypticity of B cell epitopes. *J Biol Chem* 2008;283:22737-48.
26. Khoshnoodi J, Sigmundsson K, Cartailier JP, Bondar O, Sundaramoorthy M, Hudson BG. Mechanism of chain selection in the assembly of collagen IV: a prominent role for the alpha2 chain. *J Biol Chem* 2006;281:6058-69.
27. Hellmark T, Johansson C, Wieslander J. Characterization of anti-GBM antibodies involved in Goodpasture's syndrome. *Kidney Int* 1994;46:823-9.
28. Kalluri R, Wilson CB, Weber M, et al. Identification of the alpha 3 chain of type IV collagen as the common autoantigen in antibasement membrane disease and Goodpasture syndrome. *J Am Soc Nephrol* 1995;6:1178-85.
29. Bondar O, Borza D-B, Todd P, Hudson BG. Characterization of a subpopulation of Goodpasture autoantibodies targeted to the non-collagenous domain of the alpha5(IV) chain of type IV collagen. *J Am Soc Nephrol* 2002;13:173A. abstract.
30. Zhao J, Cui Z, Yang R, Jia XY, Zhang Y, Zhao MH. Anti-glomerular basement membrane autoantibodies against different target antigens are associated with disease severity. *Kidney Int* 2009;76:1108-15.
31. Kalluri R, Melendez E, Rumpf KW, et al. Specificity of circulating and tissue-bound autoantibodies in Goodpasture syndrome. *Proc Assoc Am Physicians* 1996;108:134-9.
32. Gunwar S, Noelken ME, Hudson BG. Properties of the collagenous domain of the alpha 3(IV) chain, the Goodpasture antigen, of lens basement membrane collagen: selective cleavage of alpha (IV) chains with retention of their triple helical structure and noncollagenous domain. *J Biol Chem* 1991;266:14088-94.
33. Calvete JJ, Revert F, Blanco M, et al. Conformational diversity of the Goodpasture antigen, the noncollagenous-1 domain of the alpha3 chain of collagen IV. *Proteomics* 2006;6:Suppl 1:S237-S244.
34. Chen CR, Pichurin P, Nagayama Y, Latorra F, Rapoport B, McLachlan SM. The thyrotropin receptor autoantigen in Graves disease is the culprit as well as the victim. *J Clin Invest* 2003;111:1897-904.
35. Chazenbalk GD, Pichurin P, Chen CR, et al. Thyroid-stimulating autoantibodies in Graves disease preferentially recognize the free A subunit, not the thyrotropin holoreceptor. *J Clin Invest* 2002;110:209-17.
36. Schott M, Scherbaum WA, Morgenthaler NG. Thyrotropin receptor autoantibodies in Graves' disease. *Trends Endocrinol Metab* 2005;16:243-8.
37. Kasahara H, Matsuura E, Kaihara K, et al. Antigenic structures recognized by anti-beta2-glycoprotein I auto-antibodies. *Int Immunol* 2005;17:1533-42.
38. de Laat B, Derksen RH, van Lummel M, Pennings MT, de Groot PG. Pathogenic anti-beta2-glycoprotein I antibodies recognize domain I of beta2-glycoprotein I only after a conformational change. *Blood* 2006;107:1916-24.
39. Beck LH Jr, Bonegio RGB, Lambeau G, et al. M-type phospholipase A₂ receptor as target antigen in idiopathic membranous nephropathy. *N Engl J Med* 2009;361:11-21.

Copyright © 2010 Massachusetts Medical Society.

ELECTRONIC ACCESS TO THE JOURNAL'S CUMULATIVE INDEX

At the *Journal's* site on the World Wide Web ([NEJM.org](http://www.nejm.org)), you can search an index of all articles published since January 1975 (abstracts 1975–1992, full text 1993–present). You can search by author, key word, title, type of article, and date. The results will include the citations for the articles plus links to the full text of articles published since 1993. For nonsubscribers, time-limited access to single articles and 24-hour site access can also be ordered for a fee through the Internet ([NEJM.org](http://www.nejm.org)).