# The NEW ENGLAND JOURNAL of MEDICINE

ESTABLISHED IN 1812

JULY 2, 2009

VOL. 361 NO. 1

# M-Type Phospholipase A<sub>2</sub> Receptor as Target Antigen in Idiopathic Membranous Nephropathy

Laurence H. Beck, Jr., M.D., Ph.D., Ramon G.B. Bonegio, M.D., Gérard Lambeau, Ph.D., David M. Beck, B.A., David W. Powell, Ph.D., Timothy D. Cummins, M.S., Jon B. Klein, M.D., Ph.D., and David J. Salant, M.D.

# ABSTRACT

#### BACKGROUND

Idiopathic membranous nephropathy, a common form of the nephrotic syndrome, is an antibody-mediated autoimmune glomerular disease. Serologic diagnosis has been elusive because the target antigen is unknown.

#### METHODS

We performed Western blotting of protein extracts from normal human glomeruli with serum samples from patients with idiopathic or secondary membranous nephropathy or other proteinuric or autoimmune diseases and from normal controls. We used mass spectrometry to analyze the reactive protein bands and confirmed the identity and location of the target antigen with a monospecific antibody.

# RESULTS

Serum samples from 26 of 37 patients (70%) with idiopathic but not secondary membranous nephropathy specifically identified a 185-kD glycoprotein in nonreduced glomerular extract. Mass spectrometry of the reactive protein band detected the M-type phospholipase A<sub>2</sub> receptor (PLA<sub>2</sub>R). Reactive serum specimens recognized recombinant PLA<sub>2</sub>R and bound the same 185-kD glomerular protein as did the monospecific anti-PLA<sub>2</sub>R antibody. Anti-PLA<sub>2</sub>R autoantibodies in serum samples from patients with membranous nephropathy were mainly IgG4, the predominant immunoglobulin subclass in glomerular deposits. PLA<sub>2</sub>R was expressed in podocytes in normal human glomeruli and colocalized with IgG4 in immune deposits in glomeruli of patients with membranous nephropathy. IgG eluted from such deposits in patients with idiopathic membranous nephropathy, but not in those with lupus membranous or IgA nephropathy, recognized PLA<sub>2</sub>R.

#### CONCLUSIONS

A majority of patients with idiopathic membranous nephropathy have antibodies against a conformation-dependent epitope in  $PLA_2R$ .  $PLA_2R$  is present in normal podocytes and in immune deposits in patients with idiopathic membranous nephropathy, indicating that  $PLA_2R$  is a major antigen in this disease.

From the Boston University School of Medicine, Boston (L.H.B., R.G.B.B., D.M.B., D.J.S.); Institut de Pharmacologie Moléculaire et Cellulaire, Unité Mixte de Recherche 6097, Centre National de la Recherche Scientifique, and Université de Nice Sophia Antipolis, Valbonne, France (G.L.); and the University of Louisville School of Medicine, Louisville, KY (D.W.P., T.D.C., J.B.K.). Address reprint requests to Dr. Salant at Evans Biomedical Research Center, Rm. 504, 650 Albany St., Boston, MA 02118, or at djsalant@bu.edu.

N Engl J Med 2009;361:11-21. Copyright © 2009 Massachusetts Medical Society.

The New England Journal of Medicine

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.

DIOPATHIC MEMBRANOUS NEPHROPATHY, a common cause of the nephrotic syndrome Lin adults, is an organ-specific autoimmune disease. Despite extensive investigation, a target antigen has been elusive. Studies of membranous nephropathy in a rat model (Heymann's nephritis) established that the subepithelial immune deposits that characterize the disease are formed in situ, as a result of capping and shedding of the target antigen, megalin, from the basal surface of podocytes when it forms a complex with circulating antimegalin antibodies.1-8 Although megalin is not expressed on human podocytes, we hypothesized that a similar process, albeit with an unknown antigen, is operative in human membranous nephropathy. Additional evidence from the work of Debiec et al.9,10 supports the in situ formation of glomerular immune deposits in patients with membranous nephropathy. They described an alloimmune form of membranous nephropathy in neonates born to mothers with a deficiency of neutral endopeptidase, a protein expressed on podocytes, to which the mothers had been sensitized during previous pregnancies.

To identify the target antigen in patients with idiopathic membranous nephropathy, we used circulating antibodies from adults with this disease to detect normal glomerular proteins by using Western blotting. Subsequent analysis with the use of mass spectrometry and confirmation with the use of protein-specific reagents allowed for the identification and characterization of the predominant protein detected by these circulating antibodies.

### METHODS

# SERUM AND KIDNEY-TISSUE SAMPLES

We collected and stored serum samples from patients with membranous nephropathy, those with other glomerular or autoimmune disorders, and normal controls (healthy volunteers). The samples were assigned codes to render them anonymous. The institutional review board of the Boston University School of Medicine approved the study, and written informed consent was obtained from all the subjects. In addition, serum specimens from deceased donors with preserved renal function were obtained from the New England Organ Bank, as were kidneys, from deceased donors, that were unsuitable for transplantation. These samples were exempt from the requirement of informed consent, according to the institutional review board's approved use of organs and tissues from deceased donors for research.

Idiopathic membranous nephropathy was diagnosed by means of renal biopsy in patients who lacked features suggestive of secondary membranous nephropathy, such as antinuclear antibodies or hepatitis B antibodies in the serum (Table 1 in the Supplementary Appendix, available with the full text of this article at NEJM.org).

Glomeruli were isolated from the kidneys that were unsuitable for transplantation with the use of graded sieving,<sup>11</sup> and glomerular proteins were extracted in radioimmunoprecipitation-assay buffer (Boston BioProducts). Contaminating IgG was removed through incubation with Immobilized Protein G Plus (Fisher Scientific). Peptide N-glycosidase F (New England BioLabs) was used to deglycosylate glomerular proteins. Glomerular glycoproteins were partially purified through absorption by immobilized wheat-germ agglutinin (Vector Laboratories) and elution with N-acetyl glucosamine.

# WESTERN BLOTTING

Human glomerular extract or recombinant PLA<sub>2</sub>R expressed by human cells (see the Methods section of the Supplementary Appendix) was electrophoresed under nonreducing conditions and transferred to nitrocellulose membranes, according to standard protocols. Human serum was used as the primary antibody, at a dilution of 1:100, unless otherwise indicated. A guinea pig polyclonal antibody against the M-type phospholipase A2 receptor (hereafter referred to simply as PLA<sub>2</sub>R) was raised against full-length purified rabbit PLA<sub>2</sub>R.<sup>12</sup> Sheep antibodies against the four IgG subclasses were used as recommended by the manufacturer (Binding Site). Detecting antibodies were peroxidase-conjugated donkey antibodies against human IgG, guinea pig IgG (Jackson ImmunoResearch Laboratories), or sheep IgG (Sigma). Immunoglobulins were acid-eluted from the remnant cores of kidney-biopsy specimens from patients with membranous nephropathy, lupus membranous nephropathy, or IgA nephropathy (see the Methods section of the Supplementary Appendix). This eluted IgG was used to immunoblot the human glomerular extract or recombinant PLA<sub>2</sub>R directly.

N ENGLJ MED 361;1 NEJM.ORG JULY 2, 2009

The New England Journal of Medicine

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.

# MASS SPECTROMETRY ANALYSIS AND DATA INTERPRETATION

We excised regions of interest from gels identical to those used for Western blotting and performed in-gel tryptic digestion, as previously described.<sup>13</sup> We analyzed the resulting peptides by using a modified version of a method that couples liquid chromatography with tandem mass spectrometry<sup>14</sup> and entered the acquired data into existing databases of human proteins to identify relevant proteins in our patients.

# IMMUNOPRECIPITATION

Reactive or nonreactive human serum was incubated with human glomerular extract, immunoprecipitated with Immobilized Protein G Plus, electrophoresed in the presence of a reducing agent, and blotted with guinea pig anti-PLA<sub>2</sub>R antibodies.

### IMMUNOHISTOLOGIC ANALYSIS

Cryosections of normal human kidney and kidneybiopsy specimens obtained from patients with membranous nephropathy and made anonymous were fixed in methanol and acetone and blocked with 10% bovine serum albumin. We detected PLA<sub>2</sub>R by using guinea pig anti-PLA<sub>2</sub>R antibodies (at a dilution of 1:150), followed by a Cy3-conjugated secondary antibody (see the Supplementary Appendix), and established specificity by preabsorbing the primary antibody with a recombinant fragment of rabbit PLA,R protein<sup>15</sup> containing C-type lectinlike domains (CTLDs) 4, 5, and 6. We used a commercial rabbit antibody against human agrin at a dilution of 1:25 (Santa Cruz Biotechnology), followed by an Alexa Fluor 488conjugated secondary antibody (Invitrogen), to delineate the glomerular basement membrane. To detect podocyte nuclei, we used rabbit antibodies against Wilms' tumor 1 antigen at a 1:100 dilution (Santa Cruz Biotechnology). We detected IgG4 in immune deposits from patients with membranous nephropathy by using sheep anti-IgG4 antibodies at a dilution of 1:500 (Binding Site), followed by a fluorescein isothiocyanate-conjugated secondary antibody. Cell nuclei were visualized by including a nuclear stain (Hoechst 33342, Molecular Probes) in the mounting medium. Immunofluorescence images were obtained on a microscope (TE-2000, Nikon) linked to a Peltier-cooled CCD camera (CoolSNAP HQ, Photometrics). Confocal microscopy was performed with the use of a spinning-disk confocal instrument (UltraVIEW LCI, PerkinElmer).

# STATISTICAL ANALYSIS

Chi-square analysis was performed to compare observed with expected results. The null hypothesis was that a positive signal on Western blotting would occur in a similar proportion of subjects in each group, on the basis of chance alone.

# RESULTS

#### IDENTIFICATION OF A TARGET PROTEIN IN GLOMERULAR EXTRACTS

We performed Western blotting, under nonreducing conditions, of extracts of normal human glomeruli from donor kidneys with serum samples from 37 patients with idiopathic membranous nephropathy. A 185-kD protein band was detected in samples from 26 patients (70%) (Fig. 1). An identical protein band was seen in glomerular extracts prepared from all individual donor kidneys tested thus far. In contrast, serum from 30 normal controls, 15 patients with proteinuric conditions other than membranous nephropathy (diabetic nephropathy or focal and segmental glomerulosclerosis), and 7 patients with other autoimmune disorders did not react with this antigen when assayed under identical conditions (Fig. 1, and Table 1 in the Supplementary Appendix). Moreover, none of the serum samples from eight patients with secondary membranous nephropathy (six with lupus membranous nephropathy and two with hepatitis B-associated membranous nephropathy) were reactive with the 185-kD antigen (Fig. 1B). Chi-square analysis indicated an extremely low probability of such a distribution resulting from chance alone  $(P=2\times10^{-12})$  for the whole series; P=0.004 for secondary vs. idiopathic membranous nephropathy). Treatment with peptide N-glycosidase F caused a substantial shift in the mobility of this antigen, to approximately 145 kD, indicating that it is heavily glycosylated. All serum samples that were initially reactive with the native 185-kD band also recognized the smaller, deglycosylated band (Fig. 1A).

Both the native and N-deglycosylated forms of the antigen could be bound and substantially enriched through absorption by wheat-germ agglutinin (data not shown). The binding of the de-

13

The New England Journal of Medicine

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.



# Figure 1. Results of Western Blotting of Glomerular Proteins with Serum from Patients with Idiopathic Membranous Nephropathy.

The top of Panel A shows the results of Western blotting of extract of human glomerular proteins with serum samples from each of five patients with idiopathic membranous nephropathy (MN1 through MN5) and five patients with other proteinuric conditions (two with focal and segmental glomerulosclerosis [FGS1 and FGS2] and three with diabetic nephropathy [DN1, DN2, and DN3]). Serum samples from the five patients with membranous nephropathy all recognized a band of approximately 185 kD, whereas the samples from the patients with other diseases did not. The bottom of Panel A shows the results of Western blotting, with reactive serum samples from the five patients with membranous nephropathy, of glomerular proteins that were deglycosylated with peptide *N*-glycosidase F (PNGase F+) or not deglycosylated (PNGase F-). All five samples showed the 185-kD native antigen and a deglycosylated protein of approximately 145 kD. Panel B shows the specificity of the reactivity of serum samples to the 185-kD antigen.

> glycosylated form may reflect the presence of residual *N*-linked carbohydrates that are inaccessible to peptide *N*-glycosidase F under nonreducing conditions. Native and deglycosylated glomerular proteins were eluted from wheat-germ agglutinin and separated by means of gel electrophoresis. The regions corresponding to the 185-kD and 145-kD antigen bands on the Western blots

were excised, subjected to in-gel tryptic digestion, and analyzed by means of mass spectrometry. On the basis of the assumption that peptide sequences corresponding to the putative target antigen would be identified in both the 185-kD and 145-kD samples, we generated a working list of candidate proteins (see both Table 2 and Fig. 1 in the Supplementary Appendix). We used available antibodies and recombinant proteins to test candidate antigens and found that antibodies against PLA<sub>2</sub>R identified a glomerularprotein band of the same size as that recognized by the serum from a patient with membranous nephropathy (Fig. 2A).

# ANTI-PLA<sub>2</sub>R ANTIBODIES IN PATIENTS WITH MEMBRANOUS NEPHROPATHY

Cell-expressed recombinant human PLA<sub>2</sub>R was immunoblotted with serum from a patient with membranous nephropathy. A distinct band was detected that was slightly smaller than the corresponding band from extracts of normal human glomeruli (Fig. 2A). When both types of samples were deglycosylated, however, they migrated to the same position, suggesting a minor difference in overall glycosylation between the native protein and the recombinant form. Western-blot analysis of the same samples with the use of a monospecific polyclonal antibody against PLA<sub>2</sub>R revealed an identical pattern (Fig. 2A). The specificity of serum samples from patients with membranous nephropathy for recombinant PLA<sub>2</sub>R was established by demonstrating that they had no reactivity with extracts of cells transfected with an irrelevant plasmid (Fig. 2A in the Supplementary Appendix). All serum samples from patients with membranous nephropathy that reacted with the 185-kD glycoprotein from human glomeruli also recognized recombinant PLA<sub>2</sub>R, further suggesting that the 185-kD band from human glomeruli is indeed PLA<sub>2</sub>R.

We next confirmed that the serum samples from patients with membranous nephropathy and the anti-PLA<sub>2</sub>R antibody identified the same protein. As shown in Figure 2B (and Fig. 2B in the Supplementary Appendix), immunoprecipitates of human glomerular extract incubated with reactive serum samples from patients with membranous nephropathy yielded a 185-kD protein that was detected by anti-PLA<sub>2</sub>R antibodies on Western blotting.

N ENGLJ MED 361;1 NEJM.ORG JULY 2, 2009

The New England Journal of Medicine

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.

None of the positive serum specimens retained reactivity to the 185-kD protein when the glomerular extract was electrophoresed under reducing conditions (Fig. 3A and data not shown). Moreover, recombinant PLA<sub>2</sub>R and the native glomerular protein share the same reductionsensitive epitope (Fig. 3A). Both the monospecific anti-PLA<sub>2</sub>R antibodies and the serum samples from patients with membranous nephropathy strongly detected nonreduced recombinant and native glomerular PLA<sub>2</sub>R. Under reducing conditions, the polyclonal antibodies still detected both recombinant and native PLA<sub>2</sub>R (albeit less robustly), but the serum samples from the patients with membranous nephropathy did not. This suggests that autoantibodies in patients with membranous nephropathy uniquely recognize a conformation-dependent epitope in both the native and recombinant antigen.

The immunoglobulin profile in patients with membranous nephropathy is typical of the type 2 helper T–cell response, with the glomerular immune deposits containing predominantly IgG4subclass antibodies.<sup>16,17</sup> Similarly, serum samples from patients with membranous nephropathy are highly enriched in IgG4 autoantibodies that react with recombinant PLA<sub>2</sub>R or the native protein in glomerular extract. Although IgG4 is the predominant subclass of anti-PLA<sub>2</sub>R IgG antibodies, other subclasses are also present in smaller amounts (Fig. 3B).

# GLOMERULAR LOCATION OF PLA2R

Although we hypothesized that the target antigen resides on the surface of podocytes, we first discounted the possible formation of circulating immune complexes with soluble PLA<sub>2</sub>R. Neither serum samples from patients with membranous nephropathy nor serum samples from controls had any detectable PLA<sub>2</sub>R, even after enrichment by means of lectin binding (Fig. 3A in the Supplementary Appendix). Nor could we detect circulating immune complexes of PLA<sub>2</sub>R-IgG through either precipitation with polyethylene glycol 600018 or protein G immunoprecipitation of IgG in serum samples from either group (Fig. 3B and 3C in Supplementary Appendix). Conversely, we detected PLA<sub>2</sub>R in human glomeruli (Fig. 4A) and, more specifically, in podocytes through immunofluorescence microscopy with the monospecific anti-PLA<sub>2</sub>R antibody (Fig. 4D through 4G). PLA<sub>2</sub>R



Panel A shows the results of Western blotting of samples of human glomerular extract (HGE) and recombinant phospholipase A<sub>2</sub> receptor (rPLA<sub>2</sub>R), deglycosylated with peptide N-glycosidase F (PNGase F+) or not deglycosylated (PNGase F–), with either a reactive serum sample from a patient with membranous nephropathy (MN) or a polyclonal antibody raised against the M-type phospholipase A<sub>2</sub> receptor (PLA<sub>2</sub>R). The recombinant protein migrated to a slightly lower position than the native glomerular protein, although deglycosylation with peptide N-glycosidase F caused both the recombinant and native isoforms to migrate to the same position. Panel B shows that immunoprecipitation of native PLA<sub>2</sub>R occurred in a reactive serum sample from each of five patients with membranous nephropathy (MN1, MN2, MN3, MN6, and MN7) but not in a nonreactive sample (MN8) and not in nonreactive serum samples from normal controls (NC1 and NC2) or in the absence of serum. The immunoprecipitates were then electrophoresed under reducing conditions and subjected to Western blotting with antibodies against PLA<sub>2</sub>R (top) or against human IgG (bottom). In the reactive samples from all five patients with membranous nephropathy, PLA<sub>2</sub>R was immunoprecipitated from human glomerular extract, whereas there was no immunoprecipitation in nonreactive samples from one patient and from the controls. No PLA<sub>2</sub>R was detected when human serum was omitted from the reaction. The bottom image shows that the immunoprecipitates of samples from controls and patients with membranous nephropathy contained at least equivalent amounts of IgG.

expression in podocytes was further confirmed by positive staining with Wilms' tumor 1 antigen of PLA<sub>2</sub>R-positive cells in normal human glomeruli (Fig. 4G), as well as by the detection of PLA<sub>2</sub>R in cultured immortalized human podocytes<sup>19</sup> through the polymerase-chain-reaction assay and Western blotting (Fig. 4 in the Supple-

15

The New England Journal of Medicine

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.



# **Figure 3.** Characterization of the Antibody Response to Reduced and Nonreduced Phospholipase A<sub>2</sub> Receptor (PLA<sub>2</sub>R).

Panel A shows a Western blot in which equal amounts of human glomerular extract (HGE) and equal amounts of recombinant PLA<sub>2</sub>R (rPLA<sub>2</sub>R) were electrophoresed under reducing and nonreducing conditions. Western blotting was performed with the use of a reactive serum sample from a patient with membranous nephropathy (MN5) or a polyclonal anti-PLA<sub>2</sub>R antibody and detected with appropriate secondary antibodies. Both the serum sample from the patient with membranous nephropathy and the polyclonal anti-PLA<sub>2</sub>R antibody reacted with the native and recombinant PLA<sub>2</sub>R in the nonreduced state, but the reduced proteins were reactive with only the anti-PLA<sub>2</sub>R antibody. Panel B shows the IgG-subclass specificity to PLA2R. HGE was blotted initially with serum samples from six patients with membranous nephropathy (MN1 through MN6), followed by sheep antibodies specific for each human IgG subclass (1 through 4), and was detected with peroxidaseconjugated antisheep IgG antibody. The arrowheads indicate the fully glycosylated native PLA<sub>2</sub>R. The predominant IgG subclass that reacted with the native antigen was IgG4, with varying amounts of reactivity seen for IgG1, IgG2, and IgG3. Identical results were obtained with the use of recombinant PLA<sub>2</sub>R instead of HGE (not shown).

mentary Appendix). The specificity of PLA<sub>2</sub>R staining was shown by the near-complete inhibition of the glomerular signal after preincubation of anti-PLA<sub>2</sub>R antibody with a recombinant fragment of PLA<sub>2</sub>R containing CTLDs 4, 5, and 6 (Fig. 4B, 4E, and 4H). In contrast to podocytes, mesangial cells in normal human kidney tissue did not show staining for PLA<sub>2</sub>R (Fig. 5 in the Supplementary Appendix).

Experimental models of membranous nephropathy have shown capping and shedding of megalin from the surface of podocytes and accumulation of megalin-antimegalin antibody complexes in subepithelial immune deposits. On the basis of these data, we sought evidence that PLA<sub>2</sub>R and IgG colocalize in the glomerular immune deposits in patients with membranous nephropathy. As shown through confocal analysis of cryosections of kidney-biopsy specimens from patients with membranous nephropathy (Fig. 5A through 5F, with control samples shown in Fig. 6 and 7 in the Supplementary Appendix), PLA<sub>2</sub>R is present and colocalizes with IgG4 in a fine granular pattern that is typical of membranous nephropathy. As in the normal kidney, the staining of PLA<sub>2</sub>R could be blocked by the recombinant PLA<sub>2</sub>R fragment (Fig. 5G). Although the intensity of staining varied among the specimens from eight patients with membranous nephropathy, all revealed the same granular pattern of PLA<sub>2</sub>R (data not shown). In contrast, PLA<sub>2</sub>R and IgG4 did not colocalize in a biopsy specimen from a patient with lupus membranous nephropathy, examined under identical conditions (Fig. 5I).

To confirm that the IgG in the glomeruli from patients with membranous nephropathy was reactive with PLA<sub>2</sub>R, we eluted IgG from biopsy specimens and used it in Western blotting with native and recombinant PLA2R. IgG was successfully eluted from four biopsy samples from patients with idiopathic membranous nephropathy, one from patients with lupus membranous nephropathy, and two from patients with IgA nephropathy (Fig. 8 in the Supplementary Appendix). The IgG eluted from the samples from patients with idiopathic membranous nephropathy specifically detected the appropriately sized PLA<sub>2</sub>R bands in human glomerular extract and in cell lysates that were positive for recombinant PLA<sub>2</sub>R, whereas the IgG eluted from the three other samples from patients with immune-complex glomerular disease did not (Fig. 5J and 5K).

The New England Journal of Medicine

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.



Figure 4. Expression of the M-Type Phospholipase A<sub>2</sub> Receptor (PLA<sub>2</sub>R) in Normal Kidney Tissue and Glomeruli. Serial cryosections of the cortex of normal human kidneys were immunostained with anti-PLA<sub>2</sub>R antibody (1:150 dilution), followed by Cy3-conjugated anti-guinea pig IgG antibody (1:500 dilution). Panel A shows two positively stained glomeruli (arrowheads) and scattered tubular staining (arrow). Panel B shows an adjacent section in which the anti-PLA<sub>2</sub>R antiserum was preabsorbed with a recombinant fragment of PLA<sub>2</sub>R containing C-type lectinlike domains (CTLDs) 4, 5, and 6 (Panel H). The glomerular PLA<sub>2</sub>R staining is completely blocked by the recombinant fragment (arrowheads), whereas the scattered tubular staining (arrow) is not blocked and is therefore nonspecific. Panel C shows a human kidneytissue specimen treated with nonimmune guinea pig serum (1:200 dilution), followed by Cy3-conjugated anti-guinea pig IgG antibody, and stained with Hoechst 33342 (a nuclear stain); the inset shows a glomerulus exposed instead to anti-PLA<sub>2</sub>R antibody (1:200 dilution) but otherwise assessed under identical conditions. Next, cryosections of normal human kidney cortex were costained with antiagrin antibody, followed by an Alexa Fluor 488-conjugated antirabbit secondary antibody, to label the glomerular basement membrane (Panels D and E); in addition, PLA<sub>2</sub>R staining with guinea pig anti-PLA<sub>2</sub>R antibody was either left unblocked (Panel D) or was blocked (Panel E), as described above. The PLA<sub>2</sub>R signal is clearly present outside the glomerular basement membrane and localizes to both the cell body and the processes of the podocyte. The PLA<sub>2</sub>R staining is markedly reduced when the antibody is preabsorbed with the recombinant PLA<sub>2</sub>R fragment (Panel E). Panel F shows an enlargement of the left lower portion of the glomerulus shown in Panel D, with demarcation of the capillary lumina (asterisks) and urinary space (US) to highlight the location of the PLA<sub>2</sub>R-stained podocytes. Hoechst-stained nuclei of parietal epithelial cells in the Bowman's capsule are indicated by the arrow. Panel G shows a portion of a human glomerulus stained with anti-PLA<sub>2</sub>R antibody (as described above), anti–Wilms' tumor 1 antibody (WT1) (1:100 dilution), and Alexa Fluor 488-conjugated anti-rabbit IgG antibody (1:500 dilution) to label podocyte nuclei. The inset shows the image of the entire glomerulus from which the enlarged view was taken. Panel H illustrates the domain structure of PLA<sub>2</sub>R, which is composed of an N-terminal cysteine-rich domain (Cys-R), a fibronectin type II domain (FNII), eight CTLDs, a transmembrane domain (TM), and a short intracellular C-terminal tail (IC). The blocking fragment used in the experiments described above consists of CTLDs 4, 5, and 6 of a recombinant rabbit PLA<sub>2</sub>R.

N ENGLJ MED 361;1 NEJM.ORG JULY 2, 2009

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.





Confocal microscopic analysis of cryosections of a kidney-tissue specimen from a patient with membranous nephropathy revealed the presence of  $PLA_2R$  (Panel A) and IgG4 (Panel B), which were colocalized in the peripheral capillary walls and the glomerular basement membrane (Panel C). Panels D, E, and F are enlarged images of the boxed areas in Panels A, B, and C, respectively. The asterisks in Panel F indicate the capillary lumina. The anti-PLA<sub>2</sub>R antibody was blocked with the recombinant fragment (Panel G), with staining virtually eliminated, despite the continued presence of IgG4 (Panel H). Panel I is a confocal image of representative glomerular capillary loops in a biopsy specimen from a patient with lupus membranous nephropathy; PLA<sub>2</sub>R and IgG4 are not colocalized. IgG4 was detected with sheep anti–human IgG4 antibody (1:500 dilution) and rabbit anti–sheep IgG antibody (1:500 dilution). IgG was eluted from biopsy cores from patients with membranous nephropathy (MN), lupus membranous nephropathy (LMN), or IgA nephropathy (IgAN). This eluted IgG was used to immunoblot human glomerular extract (HGE) or recombinant PLA<sub>2</sub>R (rPLA<sub>2</sub>R) (Panel J). Only IgG eluted from the MN samples identified the native and recombinant PLA<sub>2</sub>R. Panel K shows that the IgG eluted from the MN3 biopsy sample recognized only those bands corresponding to PLA<sub>2</sub>R.

The New England Journal of Medicine

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.

#### ASSOCIATION WITH DISEASE ACTIVITY

We analyzed serum specimens collected serially from several patients in whom treatment-induced or spontaneous remission occurred. We found that, in general, autoantibodies against PLA<sub>2</sub>R were present when there was clinically significant disease activity, as measured by urinary protein and serum albumin levels (Fig. 6, and Fig. 9 in the Supplementary Appendix). In the patients with remission, there was a decline or disappearance of anti-PLA<sub>2</sub>R antibodies before the proteinuria fully resolved.

# DISCUSSION

Our findings show that PLA<sub>2</sub>R is a major target antigen in idiopathic membranous nephropathy. Seventy percent of our patients with biopsy-proven idiopathic membranous nephropathy had IgG antibodies (predominantly of the IgG4 subclass) that reacted with a reduction-sensitive epitope present on PLA<sub>2</sub>R, a glycoprotein constituent of normal human glomeruli. The distribution of PLA<sub>2</sub>R in normal glomeruli suggests that it is located on podocytes, as are megalin in rat models of membranous nephropathy and neutral endopeptidase in patients with alloimmune neonatal membranous nephropathy. PLA<sub>2</sub>R colocalizes with IgG4 in immune deposits of kidney-biopsy specimens from patients with idiopathic membranous nephropathy, and IgG eluted from such biopsy specimens is reactive with PLA<sub>2</sub>R. Moreover, serial studies in a limited number of patients suggest that the presence of circulating anti-PLA<sub>2</sub>R antibodies corresponds to disease activity. Finally, the specificity of the reactivity of anti-PLA<sub>2</sub>R antibodies in patients with idiopathic membranous nephropathy, as opposed to secondary forms of membranous nephropathy or other proteinuric diseases, suggests that the antibodies are most likely the cause, rather than a consequence, of podocyte injury and proteinuria.

There are several possible reasons why anti-PLA<sub>2</sub>R antibodies are not detected in all patients with idiopathic membranous nephropathy. First, there may be target antigens other than PLA<sub>2</sub>R; however, screening by means of Western blotting revealed no consistent alternative candidates. Second, there may be cryptic epitopes that were not revealed under our extraction and electrophoretic conditions. Third, even though all our patients had some degree of proteinuria, a proportion may



Figure 6. Antibody against the M-Type Phospholipase  $A_2$  Receptor (PLA<sub>2</sub>R) and Disease Activity in a Patient with Membranous Nephropathy.

Serum samples were collected serially from a patient with membranous nephropathy who had clinical remission after receiving immunosuppressive treatment with cyclophosphamide and prednisone (see the Clinical Vignettes section in the Supplementary Appendix). Panel A shows a decline in the urinary protein level (as measured by the protein-to-creatinine ratio) and an increase in the serum albumin level. The results of Western blotting of serum samples taken from the patient at the same times as the laboratory measurements (Panel B) show reactivity to PLA<sub>2</sub>R, either the 185-kD native form not deglycosylated by peptide *N*-glycosidase F (PNGase F–) or the 145-kD deglycosylated form (PNGase F+) in the initial serum sample (from December 2005) only. Equal loading is verified by the nonspecific detection of a 98-kD band. Equal volumes of the four serial serum samples were electrophoresed and blotted with anti–human IgG antibody (bottom), with a progressive increase in total IgG after the patient entered a period of remission from membranous nephropathy.

have no longer had immunologically active disease at the time of sample collection. Residual proteinuria in patients with membranous nephropathy is most likely due to altered podocyte architecture that results from remodeling of the glomerular basement membrane in the advanced disease state, as shown by the transplantation of kidneys from rats with active Heymann's nephritis into healthy rats.<sup>20</sup> Although proteinuria from

19

The New England Journal of Medicine

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.

the transplanted kidneys lessened in the absence of antimegalin antibodies, a moderate level of residual proteinuria persisted indefinitely in the recipient rats.

Although anti-PLA<sub>2</sub>R antibodies are predominantly of the IgG4 isotype, which corresponds to the isotype found in immune deposits in patients with idiopathic membranous nephropathy,<sup>16,21</sup> the other three IgG subclasses were variably present in serum but in smaller amounts. The presence of anti–neutral endopeptidase IgG1 and IgG4 antibodies, but not IgG4 alone, corresponds to the development of proteinuria in patients with alloimmune membranous nephropathy<sup>10</sup>; however, we could not correlate the presence or absence of anti-PLA<sub>2</sub>R IgG1 antibodies with the severity or duration of disease.

PLA<sub>2</sub>R is a type I transmembrane receptor and one of four mammalian members of the mannosereceptor family.22,23 All four have a conserved domain structure (Fig. 4H) and undergo endocytic recycling,<sup>12</sup> which may provide a constant source of accessible PLA<sub>2</sub>R at the podocyte membrane for immune-complex formation. PLA<sub>2</sub>R was initially identified as a binding protein for secreted phospholipase A2 (PLA2) and has a high binding affinity for several types of secreted PLA<sub>2</sub> in certain species.<sup>24</sup> However, human PLA<sub>2</sub>R has a different affinity for the more common types of secreted PLA<sub>2</sub>,<sup>25</sup> and its exact physiological role remains unclear.24 Human PLA2R was originally cloned from kidney tissue, in which it is expressed at high levels<sup>25</sup>; our studies suggest that its expression in human kidneys is largely confined to glomerular podocytes. In contrast, PLA<sub>2</sub>R is weakly expressed in the glomerulus in normal rats and is up-regulated in experimental models of mesangial proliferative glomerulonephritis.26 In addition, we observed that human anti-PLA<sub>2</sub>R autoantibodies do not detect PLA<sub>2</sub>R in rodent or rabbit glomerular extract (data not shown).

Anti-PLA<sub>2</sub>R autoantibodies from unrelated patients react with a similar reduction-sensitive epitope. This commonality implies that the autoantigen exists only when PLA<sub>2</sub>R is in a configuration dependent on intramolecular disulfide bonds and that the autoantibody response is restricted to this conformation. A similarly restricted immune response to a reduction-sensitive epitope has been described for Goodpasture's epitope in the noncollagenous domain NC1 in type IV collagen  $\alpha$ 3.<sup>27,28</sup> The finding of such a restricted response in patients with membranous nephropathy has interesting implications for the proximal events that initiate the immune response and limit epitope spreading to the rest of the PLA<sub>2</sub>R molecule. Since members of the mannose-receptor family exist in both extended and bent conformations that confer distinct ligand binding and multimerization capacities,<sup>29</sup> these different conformations may also play a role in exposure of the epitope in PLA<sub>2</sub>R.

The role of PLA<sub>2</sub>R in the pathogenesis of membranous nephropathy is currently unknown. The antigen may simply be a target for antibody binding and complement-mediated podocyte injury, as in experimental models of membranous nephropathy.<sup>30,31</sup> Alternatively, the antibodies may act as receptor agonists or antagonists, altering podocyte architecture and barrier function. Because PLA<sub>2</sub>R is also present in the lungs and on leukocytes,<sup>32,33</sup> it will be of interest to determine why the pathologic features are limited to the kidney.

Although there is a shorter, alternatively spliced PLA<sub>2</sub>R transcript that is predicted to encode a soluble form of the receptor,<sup>25</sup> we could not detect such a form in the serum samples from either normal controls or patients with membranous nephropathy. We also found no evidence of circulating complexes of soluble or shed PLA<sub>2</sub>R and anti-PLA<sub>2</sub>R antibodies. These findings are consistent with many earlier studies in which circulating immune complexes were not detected in patients with idiopathic membranous nephropathy<sup>3</sup> and further support the concept that immune deposits form in situ in those with the disease.

In conclusion, we identified circulating autoantibodies against PLA<sub>2</sub>R in a large proportion of patients with idiopathic membranous nephropathy whom we examined. This protein is expressed on normal glomerular podocytes and is present in the glomerular immune deposits in patients with the disease. These findings and the characteristics of PLA<sub>2</sub>R strongly suggest that it is a major target antigen in patients with idiopathic membranous nephropathy. Moreover, the specificity of these findings for idiopathic membranous nephropathy, in contrast to secondary forms of membranous nephropathy or other proteinuric kidney diseases, may have implications for the diagnosis and treatment of the disease.

The New England Journal of Medicine

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.

Supported by grants from the National Institute of Diabetes and Digestive and Kidney Diseases (DK067658 and DK30932, to Dr. Salant, and DK076743, to Dr. Powell), Amgen (to Dr. Beck), the Halpin Foundation (to Dr. Beck), Centre National de la Recherche Scientifique and Association pour la Recherche sur le Cancer (to Dr. Lambeau), and the Department of Veterans Affairs (to Dr. Klein).

Dr. Beck reports receiving grant support from Amgen and having a patent pending for a diagnostic immunoassay to detect anti- $PLA_2R$  antibodies in membranous nephropathy; Dr. Lambeau, holding patents related to the therapeutic use of secretory  $PLA_2$ proteins and their inhibitors; and Dr. Salant, receiving consulting fees from Questcor Pharmaceuticals, Cormedix, and DiObix and having a patent pending for a diagnostic immunoassay to detect anti-PLA<sub>2</sub>R antibodies in membranous nephropathy. No other potential conflict of interest relevant to this article was reported.

We thank Kathleen Dashner for help with tissue processing for immunofluorescence microscopy, Dr. Eric Boilard for help in preparing the human recombinant PLA<sub>2</sub>R and rabbit PLA<sub>2</sub>R fragment consisting of CTLDs 4 through 6, the New England Organ Bank for providing human kidneys for research purposes, the families of the organ donors, and the patients and volunteers who provided us with their serum for these studies.

#### REFERENCES

1. Van Damme BJ, Fleuren GJ, Bakker WW, Vernier RL, Hoedemaeker PJ. Experimental glomerulonephritis in the rat induced by antibodies directed against tubular antigens. V. Fixed glomerular antigens in the pathogenesis of heterologous immune complex glomerulonephritis. Lab Invest 1978;38:502-10.

2. Couser WG, Steinmuller DR, Stilmant MM, Salant DJ, Lowenstein LM. Experimental glomerulonephritis in the isolated perfused rat kidney. J Clin Invest 1978; 62:1275-87.

**3.** Couser WG, Salant DJ. In situ immune complex formation and glomerular injury. Kidney Int 1980;17:1-13.

4. Kerjaschki D, Farquhar MG. The pathogenic antigen of Heymann nephritis is a membrane glycoprotein of the renal proximal tubule brush border. Proc Natl Acad Sci U S A 1982;79:5557-61.

**5.** *Idem.* Immunocytochemical localization of the Heymann nephritis antigen (GP330) in glomerular epithelial cells of normal Lewis rats. J Exp Med 1983;157: 667-86.

**6.** Makker SP, Singh AK. Characterization of the antigen (gp600) of Heymann nephritis. Lab Invest 1984;50:287-93.

7. Raychowdhury R, Niles JL, McCluskey RT, Smith JA. Autoimmune target in Heymann nephritis is a glycoprotein with homology to the LDL receptor. Science 1989; 244:1163-5.

8. Saito A, Pietromonaco S, Loo AK, Farquhar MG. Complete cloning and sequencing of rat gp330/"megalin," a distinctive member of the low density lipoprotein receptor gene family. Proc Natl Acad Sci U S A 1994;91:9725-9.

**9.** Debiec H, Guigonis V, Mougenot B, et al. Antenatal membranous glomerulonephritis due to anti-neutral endopeptidase antibodies. N Engl J Med 2002;346:2053-60.

**10.** Debiec H, Nauta J, Coulet F, et al. Role of truncating mutations in MME gene in fetomaternal alloimmunisation and antenatal glomerulopathies. Lancet 2004;364: 1252-9.

**11.** Salant DJ, Cybulsky AV. Experimental glomerulonephritis. Methods Enzymol 1988;162:421-61.

**12.** Zvaritch E, Lambeau G, Lazdunski M. Endocytic properties of the M-type 180kDa receptor for secretory phospholipases A2. J Biol Chem 1996;271:250-7.

**13.** Powell DW, Rane MJ, Joughin BA, et al. Proteomic identification of 14-3-3zeta as a mitogen-activated protein kinase-activated protein kinase 2 substrate: role in dimer formation and ligand binding. Mol Cell Biol 2003;23:5376-87.

**14.** Powell DW, Weaver CM, Jennings JL, et al. Cluster analysis of mass spectrometry data reveals a novel component of SAGA. Mol Cell Biol 2004;24:7249-59.

15. Lambeau G, Ancian P, Barhanin J, Lazdunski M. Cloning and expression of a membrane receptor for secretory phospholipases A2. J Biol Chem 1994;269:1575-8.
16. Doi T, Mayumi M, Kanatsu K, Suehiro F, Hamashima Y. Distribution of IgG subclasses in membranous nephropathy. Clin Exp Immunol 1984;58:57-62.

**17.** Kuroki A, Iyoda M, Shibata T, Sugisaki T. Th2 cytokines increase and stimulate B cells to produce IgG4 in idiopathic membranous nephropathy. Kidney Int 2005;68: 302-10.

**18.** Brandslund I, Siersted HC, Jensenius JC, Svehag SE. Detection and quantitation of immune complexes with a rapid polyethylene glycol precipitation complement consumption method (PEG-CC). Methods Enzymol 1981;74:551-71.

**19.** Saleem MA, O'Hare MJ, Reiser J, et al. A conditionally immortalized human podocyte cell line demonstrating nephrin and podocin expression. J Am Soc Nephrol 2002;13:630-8.

**20.** Makker SP, Kanalas JJ. Course of transplanted Heymann nephritis kidney in normal host: implications for mechanism of proteinuria in membranous glomerulonephropathy. J Immunol 1989;142:3406-10.

**21.** Haas M. IgG subclass deposits in glomeruli of lupus and nonlupus membranous nephropathies. Am J Kidney Dis 1994;23:358-64.

**22**. East L, Isacke CM. The mannose receptor family. Biochim Biophys Acta 2002; 1572:364-86.

**23.** Lambeau G, Lazdunski M. Receptors for a growing family of secreted phospho-

lipases A2. Trends Pharmacol Sci 1999;20: 162-70.

**24.** Rouault M, Le Calvez C, Boilard E, et al. Recombinant production and properties of binding of the full set of mouse secreted phospholipases A2 to the mouse M-type receptor. Biochemistry 2007;46: 1647-62.

**25.** Ancian P, Lambeau G, Mattéi MG, Lazdunski M. The human 180-kDa receptor for secretory phospholipases A2: molecular cloning, identification of a secreted soluble form, expression, and chromosomal localization. J Biol Chem 1995;270: 8963-70.

**26.** Beck S, Beck G, Ostendorf T, et al. Upregulation of group IB secreted phospholipase A(2) and its M-type receptor in rat ANTI-THY-1 glomerulonephritis. Kidney Int 2006;70:1251-60.

**27.** Hellmark T, Burkhardt H, Wieslander J. Goodpasture disease: characterization of a single conformational epitope as the target of pathogenic autoantibodies. J Biol Chem 1999;274:25862-8.

**28.** Hudson BG. The molecular basis of Goodpasture and Alport syndromes: beacons for the discovery of the collagen IV family. J Am Soc Nephrol 2004;15:2514-27.

**29.** Llorca O. Extended and bent conformations of the mannose receptor family. Cell Mol Life Sci 2008;65:1302-10.

**30.** Nangaku M, Shankland SJ, Couser WG. Cellular response to injury in membranous nephropathy. J Am Soc Nephrol 2005;16:1195-204.

**31.** Cybulsky AV, Quigg RJ, Salant DJ. Experimental membranous nephropathy redux. Am J Physiol Renal Physiol 2005;289: F660-F671.

**32.** Granata F, Petraroli A, Boilard E, et al. Activation of cytokine production by secreted phospholipase A2 in human lung macrophages expressing the M-type receptor. J Immunol 2005;174:464-74.

**33.** Silliman CC, Moore EE, Zallen G, et al. Presence of the M-type sPLA(2) receptor on neutrophils and its role in elastase release and adhesion. Am J Physiol Cell Physiol 2002;283:C1102-C1113.

Copyright © 2009 Massachusetts Medical Society.

N ENGLJ MED 361;1 NEJM.ORG JULY 2, 2009

Downloaded from nejm.org at UNIV OF LOUISVILLE on May 22, 2012. For personal use only. No other uses without permission.