

GENVOYA

**ELVITEGRAVIR 150MG / COBICISTAT 150MG /
EMTRICITABINE 200MG / TENOFOVIR
ALAFENAMIDE 10MG**



One pill, once a day.

Genvoya is a prescription pill that contains 4 medications in one tablet and is approved by the FDA for the treatment of HIV infection in adults and children over 12 years of age as a complete regimen.

How should I take Genvoya?

- Take Genvoya **with food** at the **same time everyday**

What drugs interact with Genvoya?

- Genvoya should be taken **2 hours before or 6 hours after** antacids (such as Tums®, Mylanta®, Maalox®).
- Genvoya interacts with **A LOT** of other medications – be sure all of your health care providers know you are on this medicine.
- If you are unsure if a medicine is safe to take with your Genvoya, call the clinic pharmacist to double check.

What should I do if I forget a dose?

- If you miss a dose, take the missed dose as soon as you remember!
- If it is within 4 hours of your next dose, skip the missed dose and take the next dose at your regular time.
- Do NOT take two doses at the same time.

Resources: Gilead Important Facts, Genvoya; Department for Health and Human Services, AidsInfo, Genvoya.

Side Effects

The most common side effects of Genvoya are nausea and headache.

Most side effects go away within a few days of starting your medicine

Contact the clinic if any of these side effects become severe or prevent you from taking your medicine.

Contact the clinic if you experience any of the following:

- Unusual muscle pain
- Dark colored urine
- Rash

One pill, once a day with food

