

ODEFSEY

**RILPIVIRINE 25MG / EMTRICITABINE 200MG /
TENOFIVIR ALAFENAMIDE 25MG**



One pill, once a day.

Odefsey is a prescription pill that contains 3 medications in one tablet and is approved by the FDA for the treatment of HIV infection in adults and children over 12 years of age as a complete regimen.

How should I take Odefsey?

- Take Odefsey **with food** at the **same time everyday**

What drugs interact with Odefsey?

- Odefsey **cannot** be taken with proton pump inhibitors, such as omeprazole (Protonix®), esomeprazole (Nexium®), lansoprazole (Prevacid®), as well as others.
- Odefsey has to be separated from other antacids and heartburn medicines (such as Tums®, Mylanta®, Maalox®, Zantac®, or Pepcid®).
- If you are unsure if a medicine is safe to take with your Odefsey, call the clinic pharmacist to double check.

What should I do if I forget a dose?

- If you miss a dose, take the missed dose as soon as you remember!
- If it is within 4 hours of your next dose, skip the missed dose and take the next dose at your regular time.
- Do NOT take two doses at the same time.

Side Effects

The most common side effects of Odefsey are **nausea, depression, and drowsiness.**

Most side effects go away within a few days of starting your medicine

Contact the clinic if any of these side effects become severe or prevent you from taking your medicine.

Contact the clinic if you experience any of the following:

- Depression
- Change in the color or amount of urine passed
- Rash

One pill, once a day with food

