

# PREZCOBIX

**DARUNAVIR 800 MG / COBICISTAT 150 MG**



**Once a day, as part of a complete regimen**

PrezcoBix is a prescription pill that contains 2 medications in one tablet and is approved by the FDA for the treatment of HIV infection in adults and children over 12 years in combination with other medications.

PrezcoBix must be taken in combination with other HIV medications.

## How should I take PrezcoBix?

- Take PrezcoBix **with food** at the **same time everyday**
- **Always** take PrezcoBix **in combination with** other HIV medications

## What drugs interact with PrezcoBix?

- PrezcoBix interacts with **A LOT** of other medications – be sure all of your health care providers know you are on this medicine.
- If you are unsure if a medicine is safe to take with your PrezcoBix, call the clinic pharmacist to double check.

## What should I do if I forget a dose?

- If you miss a dose, take the missed dose as soon as you remember!
- If it is within 4 hours of your next dose, skip the missed dose and take the next dose at your regular time.
- Do **NOT** take two doses at the same time.

## Side Effects

The most common side effects of PrezcoBix are nausea, diarrhea, and rash.

**Most side effects go away within a few days of starting your medicine**

**Contact the clinic if any of these side effects become severe or prevent you from taking your medicine.**

**Contact the clinic or go to the emergency department if you experience any of the following:**

- **Rash**
- **Signs or symptoms of allergic reaction**



**One pill daily**