

TIVICAY

DOLUTEGRAVIR 50 MG



Once a day, as part of a complete regimen

Tivicay is a prescription pill that is approved by the FDA for the treatment of HIV infection in adults and children over 12 years of age in combination with other medications.

Tivicay must be taken in combination with other HIV medications.

How should I take Tivicay?

- Take Tivicay **with your other HIV medications** at the **same time everyday**.
- Tivicay can be taken with or without food.

What drugs interact with Tivicay?

- Tivicay should be taken **2 hours before or 6 hours after** antacids, iron/calcium supplements, multivitamins or prenatal vitamins, or any other medications containing aluminum, calcium, magnesium, or sucralfate.
- If you are unsure if a medicine is safe to take with your Tivicay, call the clinic pharmacist to double check.

What should I do if I forget a dose?

- If you miss a dose, take the missed dose as soon as you remember!
- If it is within 4 hours of your next dose, skip the missed dose and take the next dose at your regular time.
- Do NOT take two doses at the same time.

Resources: GlaxoSmithKline, Tivicay; Department for Health and Human Services, AidsInfo, Tivicay.

Side Effects

The most common side effects of Tivicay are nausea and difficulty sleeping

Most side effects go away within a few days of starting your medicine

Contact the clinic if any of these side effects become severe or prevent you from taking your medicine.

Contact the clinic if you experience any of the following:

- Unusual muscle pain
- Dark colored urine
- Rash



Once pill daily

with other HIV medications